

DISTRIBUTIONAL RESULTS FOR THE SHORTEST DISTANCE BETWEEN TRAJECTORIES OF DIFFERENT DYNAMICS

ROMAIN AIMINO, THÉOPHILE CABY, JORGE MILHAZES FREITAS, AND DUARTE SÁ PINHO

ABSTRACT. We establish Extreme Value Distributions for the closest encounter between trajectories generated by different maps defined in the same reference phase space. For a class of strongly mixing maps, we show that the limit distribution depends on the length of the different trajectories and the co-dimension of the associated invariant measures. It is also modulated by an Extremal Index, that informs on the tendency of nearby points to diverge along with the evolution of their respective dynamics, serving as an indicator of their compatibility. We give a formula for this quantity for a class of chaotic maps of the interval and for the co-dimension in the case when the respective measures admit densities with isolated zeros and singularities. We present diverse examples of systems satisfying these assumptions and compute the different parameters modulating the limit distribution.

1. INTRODUCTION

The statistical behaviour of the shortest distance between orbits and trajectories of dynamical systems has been studied for several decades, beginning at least with the pioneering work of [15] and its subsequent developments in [14]. In recent years, however, the subject has attracted renewed attention, partly due to its connection with sequence matching in symbolic dynamics, observed in [6].

Distributional results for trajectories generated by the same dynamical system were obtained in [19, 10] using techniques from extreme value theory, while almost sure (a.s.) convergence results for the shortest distance between two orbits appear in [6], and for multiple orbits in [7]. These a.s. results were later generalised to observed systems [16, 9] and to slowly mixing systems [32]. These a.s. developments were carried further in the work [34], which considers two trajectories driven by different maps and derives asymptotic results for the shortest distance, while the observed counterpart for distinct dynamical systems was recently investigated in [8].

2010 *Mathematics Subject Classification.* 37A25, 37A50, 37B20, 37C30, 60G70.

Key words and phrases. Synchronisation of orbits, shortest distance, extreme values, spectral methods, extremal index.

All authors were partially supported by CMUP, member of LASI, which is financed by national funds through FCT – Fundação para a Ciência e a Tecnologia, I.P., under the projects with reference UID/00144/2025 and associated DOI given by <https://doi.org/10.54499/UID/00144/2025>. DSP gratefully acknowledges the scholarship awarded by the Calouste Gulbenkian Foundation under the Novos Talentos programme, during which this project was initiated. TC was supported by ISblue project, Interdisciplinary graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French government under the program 'Investissements d'Avenir' embedded in France 2030.

Further developments include the study of shortest distances between partial orbits [31] and extensions to conformal iterated function systems [35]. Very recent advances on the minimal distance between random orbits were obtained in [27].

In this work, we study multiple trajectories generated by possibly distinct maps and establish full distributional results for their minimal mutual distance. In this setting, an extreme event corresponds to a strong form of synchronisation: two or more trajectories, starting from independent initial conditions and evolving under different dynamics, approach each other arbitrarily close at the same time. We derive the limiting distribution, identify the associated modulating parameters — namely the co-dimension C_q and the Extremal Index θ — and provide several explicit examples illustrating the theory.

The co-dimension plays an important role in the normalising sequences that assure distributional convergence. It essentially accounts for the irregularities of the invariant measure, namely, when zeros or singularities of the density arise with a significant impact or when its support is made out of fractal sets, for example. In the case of smooth invariant densities the co-dimension is 1.

The Extremal Index which changes the limiting law is seen here to serve as an indicator of the compatibility between the factor dynamics. Essentially, it detects when the factor dynamics have a tendency to remain synchronised, once a synchronisation occurs. Typically, different dynamics have irrelevant compatibility, in the sense that most of the same initial conditions lead to different trajectories (in measure-theoretic terms), which is reflected in an Extremal Index equal to 1. This is because, in these extreme value approaches, as in [19, 18], we consider an observable function that takes maximal values when the orbits synchronise and then, since when the trajectories synchronise there is no strong propensity for the system to remain synchronised, then we have no clustering of high observations and the Extremal Index takes the value 1. By contrast, when the factor dynamics coincide, namely, when one studies trajectories generated by the same map, as in [15, 14, 19, 10], synchronisation persists indefinitely once it occurs. In particular, this means that when we have near-synchronisation, corresponding to the observation of a high value, there is a clear predisposition to remain close to synchronisation, which creates clustering of high observations, reflected in an Extremal Index less than 1. Then, as observed in [15], the Extremal Index can be interpreted as an average of the rate of expansion of the factor dynamics. In more physical terms, the Extremal Index reflects the positive Lyapunov exponent, in low dimensional dynamics and the metric entropy in higher dimensions, as insightfully pointed out in [19, 11]. This is because these quantities essentially express the ‘tendency’ of the orbits to move away from each other.

In this work, we introduce examples in which the factor dynamics are compatible without being identical. More precisely, the dynamics coincide on part of the phase space and differ on its complement. In this setting, the Extremal Index becomes a weighted average of the two regimes: it captures the average expansion rate on the region where the dynamics agree, while taking the value 1 on the complementary region. The corresponding weights are given precisely by the relative measures of these two parts of the phase space.

We will use spectral methods of perturbed Perron-Frobenius operators introduced by Keller and Liverani in [29, 30] to establish an abstract result which will enable us to apply their framework to the problem of obtaining distributional limits for the closest distance between trajectories generated by possibly distinct dynamics. Then, we will develop applications of

this result to illustrate the roles of the co-dimension and, especially, the Extremal Index in the interpretation of the different regimes.

The structure of the paper is as follows. In Section 2, we establish the setting and formulate the problem to be studied. In Section 3, we revise Keller and Liverani's framework, introduce the notation and gather the main tools. In Section 4, we prove the abstract result that we will use subsequently to study the distributional limits mentioned earlier. In Section 5, we provide general conditions that allow us to prove that the Extremal Index is less than 1, corresponding to the typical case, when the factor dynamics is incompatible, which means that we have unclustered synchronisation profiles. In Section 6, we show the usefulness of the Extremal Index as an indicator of the compatibility of the factor dynamics, using the developed tools to compute it for linear dynamical systems, with which we play to provide an example of distinct factor systems with a compatible component. In Section 7, we provide further applications of the theory developed to handle factor dynamics such as perturbed linear maps. We end with an appendix where we discuss the possible role the co-dimension and its impact on the estimation of the quantities considered earlier.

2. THE SETTING

Let $\mathcal{M}$ be a compact metric space equipped with a distance d . For some $q \in \mathbb{N}$, $q \geq 2$, we consider the dynamical systems $(\mathcal{M}, T_1, \mu_1), (\mathcal{M}, T_2, \mu_2), \dots, (\mathcal{M}, T_q, \mu_q)$, where for each $i = 1, \dots, q$, $T_i : \mathcal{M} \rightarrow \mathcal{M}$ is a discrete transformation that leaves the probability measure μ_i invariant. We will denote $\bar{\mu} := \mu_1 \times \mu_2 \times \dots \times \mu_q$ the product measure with support in $\mathcal{M}^q$, and $\bar{T} : \mathcal{M}^q \rightarrow \mathcal{M}^q$ the product map

$$\bar{T}(\bar{x}) := (T_1(x_1), \dots, T_q(x_q)), \quad (2.1)$$

where $\bar{x} = (x_1, \dots, x_q) \in \mathcal{M}^q$.

Definition 2.1. We call the *co-dimension* of the measures $\mu_1, \dots, \mu_q$ the following quantity (if it exists):

$$C_q(\mu_1, \dots, \mu_q) := \lim_{r \rightarrow 0} \frac{\log \int_{\mathcal{M}} \prod_{i=2}^q \mu_i(B_r(x)) d\mu_1(x)}{(q-1) \log r}, \quad (2.2)$$

where $B_r(x)$ denotes the ball of radius r centred at $x \in \mathcal{M}$.

Remark 2.2. By Fubini, we have that C_q is symmetric in $\mu_1, \dots, \mu_q$. Moreover, when $\mu_1 = \mu_2 = \dots = \mu_q$, then $C_q(\mu_1, \dots, \mu_q) = D_q(\mu_1)$, which stands for the generalised Rényi dimension of order q of μ_1 .

We define the observable function

$$\begin{aligned} \varphi : [0, 1]^q &\longrightarrow \mathbb{R} \cup \{\infty\} \\ (x_1, \dots, x_q) &\longmapsto -\log \max_{j \in \{2, \dots, q\}} d(x_1, x_j), \end{aligned} \quad (2.3)$$

which will serve as a synchronisation indicator. We define now the process $(Y_i)_i$ simply by evaluating φ along the orbits of the system:

$$Y_i(\bar{x}) := \varphi \circ \bar{T}^i(\bar{x}) = -\log \max_{j=2, \dots, q} d(T_1^i(x_1), T_j^i(x_j)),$$

where $\bar{x} \in \mathcal{M}^q$ is drawn from $\bar{\mu}$.

Note that $Y_i = \infty$ means that the system is synchronised at time i as all factors are in the same state, *i.e.*, $T_1^i(x_1) = T_2^i(x_2) = \dots = T_q^i(x_q)$.

As usual, in order to study the Extreme Value properties of this process, we take a sequence of thresholds $(u_n(s))_n$, $s > 0$, satisfying

$$\bar{\mu}(Y_0 > u_n(s)) \sim \frac{s}{n}. \quad (2.4)$$

Since the q trajectories are independent (chosen with respect to the product measure),

$$\bar{\mu}(Y_0 > u_n(s)) = \int_{\mathcal{M}} \prod_{i=2}^q \mu_i(B_{r_n}(x)) d\mu_1(x) \sim e^{-u_n(s)C_q(q-1)}, \quad (2.5)$$

where $r_n = e^{-u_n}$. Matching (2.4) and (2.5) gives

$$u_n(s) = \frac{\log n}{C_q(q-1)} - \frac{\log s}{C_q(q-1)}.$$

The quantity $\bar{\mu}(Y_0 > u_n)$ corresponds to the probability that all points $x_1, \dots, x_q$ lie in the ball $B_{r_n}(x_1)$, *i.e.*, that the product dynamics enters the target set

$$\Delta_n = \left\{ \bar{x} \in \mathcal{M}^q : \max_{j=2, \dots, q} d(x_1, x_j) < r_n \right\}. \quad (2.6)$$

Following [22], we define

$$M_n := \max\{Y_0, \dots, Y_{n-1}\}. \quad (2.7)$$

Observe that the event $\{M_n \leq u_n\}$ corresponds to the dynamics never entering Δ_n , during the first n steps, which means that the systems never came close to being synchronised. Also note that the value of M_n is related to the shortest distance reached by q trajectories up to time n .

3. A SPECTRAL APPROACH TO STUDY EXTREME VALUE LAWS

In [29], Keller and Liverani introduced a powerful framework of spectral perturbation methods, which, later, they developed further and applied to the study Rare Events and Escape Rates in [30]. This framework has since then been exploited to study Extreme Values in [28, 5, 3, 4], for example. We will use these spectral perturbation techniques to obtain our main results and, therefore, we start by describing them and recalling their main features.

3.1. Abstract perturbation results. Let $(V, \|\cdot\|)$ be a real or complex normed vector space with dual $(V^*, \|\cdot\|)$. Suppose that we have a family of operators $(P_\varepsilon)_{\varepsilon \in E}$, satisfying

$$\|P_\varepsilon\| \leq M, \quad (A0)$$

where $E \subseteq \mathbb{R}$ is a closed set of parameters having $\varepsilon = 0$ as an accumulation point. Suppose that there are $\lambda_\varepsilon \in \mathbb{C}$, $\varphi_\varepsilon \in V$, $\nu_\varepsilon \in V^*$ and linear operators $Q_\varepsilon : V \rightarrow V$ such that

$$\lambda_\varepsilon^{-1} P_\varepsilon = \varphi_\varepsilon \otimes \nu_\varepsilon + Q_\varepsilon \quad (A1)$$

$$P_\varepsilon(\varphi_\varepsilon) = \lambda_\varepsilon \varphi_\varepsilon, \nu_\varepsilon P_\varepsilon = \lambda_\varepsilon \nu_\varepsilon, Q_\varepsilon(\varphi_\varepsilon) = 0, \nu_\varepsilon Q_\varepsilon = 0 \quad (A2)$$

$$\sum_{n=0}^{\infty} \sup_{\varepsilon \in E} \|Q_\varepsilon^n\| =: C_1 < \infty. \quad (A3)$$

Conditions (A1)+(A2) imply that $\nu_\varepsilon(\varphi_\varepsilon) = 1$, for all $\varepsilon \in E$. We also control the size of φ_ε relatively to the size of φ_0 .

$$\nu_0(\varphi_\varepsilon) = 1 \text{ and } \sup_{\varepsilon \in E} \|\varphi_\varepsilon\| =: C_2 < \infty. \quad (\text{A4})$$

Define

$$\delta_\varepsilon := \nu_0((P_0 - P_\varepsilon)(\varphi_0)) \quad (3.1)$$

and suppose that there exists $C_3 > 0$ such that

$$\eta_\varepsilon := \|\nu_0(P_0 - P_\varepsilon)\| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0 \quad (\text{A5})$$

$$\eta_\varepsilon \cdot \|(P_0 - P_\varepsilon)(\varphi_0)\| \leq C_3 |\delta_\varepsilon|. \quad (\text{A6})$$

The main abstract result in [30, (Theorem 2.1)] tells us that under assumptions (A0)-(A6), then, one of the following holds

- (a) There is $\varepsilon_0 > 0$ such that $\lambda_\varepsilon = \lambda_0$ if $\varepsilon \leq \varepsilon_0$ and $\delta_\varepsilon = 0$.
- (b) If $\delta_\varepsilon \neq 0$ for all sufficiently small $\varepsilon \in E$ and if

$$p_k := \lim_{\varepsilon \rightarrow 0} p_{k,\varepsilon} := \lim_{\varepsilon \rightarrow 0} \frac{\nu_0((P_0 - P_\varepsilon) P_\varepsilon^k (P_0 - P_\varepsilon)(\varphi_0))}{\delta_\varepsilon} \quad (\text{A7})$$

exists for each integer $k \geq 0$, then

$$\lim_{\varepsilon \rightarrow 0} \frac{\lambda_0 - \lambda_\varepsilon}{\delta_\varepsilon} = 1 - \sum_{k=0}^{\infty} \lambda_0^{-(k+1)} p_k. \quad (3.2)$$

Moreover, as seen in [29], the following conditions are sufficient to verify (A0)-(A4):

(B1) The operator P_0 has the form

$$P_0 = \varphi_0 \otimes \nu_0 + Q_0, \text{ where } \nu_0(\varphi_0) = 1, \nu_0 Q_0 = 0, Q_0 \varphi_0 = 0, \quad (\text{B1.1})$$

and the spectral radius of Q_0 is strictly less than 1.

(B2) There exists $r \in (0, 1)$, $D > 0$ and a semi-norm $|\cdot|_w \leq \|\cdot\|$ in V such that

$$\text{The residual spectrum of } P_\varepsilon \text{ is contained in } \{z \in \mathbb{C} : |z| \leq r\}. \quad (\text{B2.1})$$

$$\forall \varepsilon \in E, \forall f \in V, \forall n \in \mathbb{N}, |P_\varepsilon^n f|_w \leq D |f|_w \quad (\text{B2.2})$$

$$\forall \varepsilon \in E, \forall f \in V, \forall n \in \mathbb{N}, \|P_\varepsilon^n f\| \leq D r^n \|f\| + D |f|_w \quad (\text{B2.3})$$

(B3) There exists a decreasing upper-semicontinuous function $\pi_\varepsilon : E \rightarrow (0, \infty)$ such that

$$\lim_{\varepsilon \rightarrow 0} \pi_\varepsilon = 0 \text{ and } \forall f \in V, \forall \varepsilon \in E : |P_\varepsilon f - P_0 f|_w \leq \pi_\varepsilon \|f\|. \quad (\text{B3.1})$$

Remark 3.1. The following notation will be useful. Let $R : V \rightarrow V$ be a linear operator. We define

$$\|R\| := \sup \{|Rf|_w : f \in V, \|f\| \leq 1\}. \quad (3.3)$$

With this notation, we can write the last part of condition (B3) as $\|P_\varepsilon - P_0\| \leq \pi_\varepsilon$.

3.2. Extreme Values from a spectral approach. Following [28], we consider the so called Rare event Perron–Frobenius operators (REPFO) setting. Let $[0, 1]^q$ be the phase space, and Leb the reference measure defined on the Borel subsets of $[0, 1]^q$. Assume that V is a Banach space of functions continuously embedded and dense in $L^1([0, 1]^q, \text{Leb})$ which contains the constant functions and P_0 is the transfer operator of a non-singular map $T : [0, 1]^q \rightarrow [0, 1]^q$ with respect to the reference measure Leb . Recall that in this setting, Leb defines a functional on V , by associating to each $f \in V$ the number $\int f d\text{Leb}$. We assume that $\nu_0 = \text{Leb}$. We also assume that there exists a family $(A_n)_{n \in \mathbb{N}}$ of sufficiently regular subsets of $[0, 1]^q$ such that

- (R1) For all $f \in V$, $\mathbb{1}_{A_n} \cdot f \in V$.
- (R2) The operators P_n , defined by $P_n f = P_0(\mathbb{1}_{[0, 1]^q \setminus A_n} f)$ satisfy assumptions (B1)-(B3), with $|\cdot|_w = \|\cdot\|_{L^1([0, 1]^q, \text{Leb})}$.
- (R3) There exists $C > 0$ such that for all $f \in V$, $|\nu_0(\mathbb{1}_{A_n} f)| \cdot \|\mathbb{1}_{A_n} \varphi_0\| \leq C \|f\| \cdot |\nu_0(\mathbb{1}_{A_n} \varphi_0)|$.

We note that since (B1) is satisfied, $\mu := \varphi_0 \text{Leb}$ is the unique invariant measure that is absolutely continuous with respect to the reference measure Leb . In what follows, we will also denote $h = \varphi_0$.

Lemma 3.2. *Assume that V is a space of functions on $[0, 1]^q$ that is a Banach algebra and is continuously embedded in $L^\infty([0, 1]^q, \text{Leb})$. Suppose that $\inf_{x \in \bigcup_n A_n} h > 0$. In addition, suppose that $\|\mathbb{1}_{A_n}\|$ is uniformly bounded. Then (R3) is satisfied.*

Proof. Let $f \in V$. Since $V \subseteq L^\infty([0, 1]^q, \text{Leb})$,

$$|\nu_0(\mathbb{1}_{A_n} f)| = \left| \int \mathbb{1}_{A_n} f d\text{Leb} \right| \leq \|f\|_{L^\infty} \cdot \left| \int \mathbb{1}_{A_n} d\text{Leb} \right|.$$

Using the fact that the injection of V in L^∞ is continuous, we get

$$|\nu_0(\mathbb{1}_{A_n} f)| \leq C_1 \|f\| \cdot \left| \int \mathbb{1}_{A_n} d\text{Leb} \right|.$$

The assumption that $\inf_{x \in \bigcup_n A_n} h > 0$ allows us to evaluate $\mathbb{1}_{A_n}$ with respect to the measure $\mu := h \text{Leb}$:

$$\left| \int \mathbb{1}_{A_n} d\text{Leb} \right| \leq \frac{1}{\inf_{x \in \bigcup_n A_n} h} \left| \int \mathbb{1}_{A_n} h d\text{Leb} \right|.$$

Since $(V, \|\cdot\|)$ is a Banach algebra and $\|\mathbb{1}_{A_n}\|$ is uniformly bounded, we conclude that

$$\|\mathbb{1}_{A_n} h\| \leq C'_2 \|\mathbb{1}_{A_n}\| \cdot \|h\| \leq C_2.$$

Putting all of this together,

$$|\nu_0(\mathbb{1}_{A_n} f)| \cdot \|\mathbb{1}_{A_n} h\| \leq \frac{C_1 C_2 \|f\|}{\inf_{x \in \bigcup_n A_n} h} \left| \int \mathbb{1}_{A_n} h d\text{Leb} \right| \leq C \|f\| \cdot |\nu_0(\mathbb{1}_{A_n} h)|,$$

which concludes the proof. $\square$

Now, we illustrate how one can establish the existence of a limiting EVL in a REPFO setting. According to [28, Proposition 1], the operators $(P_n)_n$ satisfy conditions (B1)-(B3), so

$$P_n = \varphi_n \otimes \nu_n + Q_n \quad \text{and} \quad \sup_n \|Q_n^k\| \leq K_3 r^k, \quad \text{where} \quad 0 < r < 1.$$

Letting $\delta_n := \int \mathbb{1}_{A_n} h d\text{Leb} = \mu(A_n) \neq 0$, for all sufficiently large n , we have by (3.2) the expansion:

$$\lim_{n \rightarrow \infty} \frac{1 - \lambda_n}{\delta_n} = \theta := 1 - \sum_{j=0}^{\infty} p_j, \quad (3.4)$$

if we assume now that for each $k \in \mathbb{N}$, the following limit exists:

$$p_k = \lim_{n \rightarrow \infty} p_{k,n} := \frac{\mu(A_n \cap T^{-1}A_n^c \cap \dots \cap T^{-k}A_n^c \cap T^{-(k+1)}A_n)}{\mu(A_n)}. \quad (3.5)$$

The first hitting time to the set A_n is $\tau_n(x) = \inf\{i \geq 0 : T^i x \in A_n\}$. Since $\nu_0 = \text{Leb}$ and that $\nu_0(\varphi_n) = 1$, we have

$$\begin{aligned} \mu(\{\tau_n \geq k\}) &= \int_{\{\tau_n \geq k\}} h d\text{Leb} = \int \prod_{i=0}^{k-1} (\mathbb{1}_{[0,1]^q \setminus A_n} \circ T^i) h d\text{Leb} \\ &= \int P_n^k h d\text{Leb} \\ &= \lambda_n^k \nu_n(h) + \mathcal{O}(\lambda_n^k \|Q_n^k\| \cdot \|h\|). \end{aligned}$$

The quantity $\nu_n(h) \rightarrow 1$ as $n \rightarrow \infty$. Suppose that $n\mu(A_n) \rightarrow s$ for some $s > 0$. Then,

$$\lim_{n \rightarrow \infty} \mu(\{\tau_n \geq n\}) = \lim_{n \rightarrow \infty} \exp(-n\mu(A_n)\theta + o(\mu(A_n)) \cdot n) = e^{-\theta s}. \quad (3.6)$$

In particular, returning to our original synchronisation problem described in Section 2, if $\varphi : [0, 1]^q \rightarrow \mathbb{R} \cup \{\infty\}$ is given as in (2.3) and there exists a sequence u_n such that $n\mu(\{\varphi > u_n\}) \rightarrow s$ and if the sets $\Delta_n = \{\varphi > u_n\}$ satisfy (R1)-(R3), then

$$\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-\theta s}.$$

4. ABSTRACT LIMITING RESULTS FOR THE PRODUCT OF INTERVAL MAPS

In this section we prove the existence of an extreme value law for the process (Y_i) of Section 3 using the spectral approach. We consider $\mathcal{M} = [0, 1]^q$ and denote by $\mu_i = h_i \text{Leb}$ the absolutely continuous invariant probability measures. We assume that the product map $\tilde{T} : [0, 1]^q \rightarrow [0, 1]^q$ is piecewise smooth with branches indexed by j to distinguish from the indices $i = 1, \dots, q$ of the factors; we write $T_{j,i}$ for the restriction of the factor T_i of $\tilde{T}$ to the domain of injectivity of the j -th branch of $\tilde{T}$. For this kind of maps the transfer operator can be expressed as

$$P(f)(\mathbf{x}) = \sum_{\tilde{T}(\mathbf{y})=\mathbf{x}} \frac{f(\mathbf{y})}{\text{Jac } \tilde{T}(\mathbf{y})}.$$

4.1. Quasi-Hölder space and Lasota-Yorke inequality. Before stating our main result, we need to introduce the functional space where we will work on. It was introduced by Saussol in [33]. We define the quantity

$$|f|_\alpha = \sup_{0 < \epsilon \leq \epsilon_0} \epsilon^{-\alpha} \int_{[0,1]^q} \text{osc}(f, B_\epsilon(x)) dx. \quad (4.1)$$

and the α -Quasi-Hölder space by

$$V_\alpha = \{f \in L^1([0, 1]^q, \text{Leb}) : |f|_\alpha < \infty\}. \quad (4.2)$$

The norm considered in V_α is $\|\cdot\|_\alpha = \|\cdot\|_{L^1([0, 1]^q, \text{Leb})} + |\cdot|_\alpha$. As shown in [33], V_α is a Banach space, compactly embedded in L^1 , continuously embedded in L^∞ , and is a Banach algebra. The space V_α is independent of the choice of the parameter ϵ_0 , and different choices of ϵ_0 yield equivalent norms.

We will assume that the transfer operator of the map $\tilde{T}$ satisfies a Lasota-Yorke inequality for the first iteration:

(LY) There exist $\eta \in (0, 1)$ and $D > 0$ such that, for all $f \in V_\alpha$,

$$\|Pf\|_\alpha \leq \eta\|f\|_\alpha + D\|f\|_{L^1}. \quad (\text{LY})$$

4.2. Synchronisation distributional limits. We now state and prove the main result of this section. Consider the set $[0, 1]^q$, where q is an integer greater or equal to 2. Let Leb be the reference measure. Consider the product dynamical system $\bar{T}: [0, 1]^q \rightarrow [0, 1]^q$ and the synchronisation observable function $\varphi: [0, 1]^q \rightarrow \mathbb{R} \cup \{\infty\}$ given in (2.1) and (2.3), respectively.

Theorem 4.1. *Assume that the transfer operator of $\bar{T}$ satisfies conditions LY and (B1) and that h is bounded away from zero, i.e., there exists $c > 0$ such that $h \geq c$. Let $\mu := h \text{Leb}$. Assume that there exists a sequence $(u_n)_n$ such that*

$$\lim_{n \rightarrow \infty} n\mu(\Delta_n) = \lim_{n \rightarrow \infty} n\mu(\{\varphi > u_n\}) = s, \quad (4.3)$$

for some $s > 0$, and, moreover, assume that for each $k \geq 0$, the limit

$$p_k := \lim_{n \rightarrow \infty} \frac{\mu(\Delta_n \cap \bar{T}^{-1}\Delta_n^c \cap \dots \cap \bar{T}^{-k}\Delta_n^c \cap \bar{T}^{-(k+1)}\Delta_n)}{\mu(\Delta_n)} \quad (4.4)$$

exists. Then, for all $s > 0$, we have

$$\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-\theta s}, \quad (4.5)$$

where $\theta := 1 - \sum_{k=0}^{\infty} p_k$.

Remark 4.2. This theorem guarantees the existence of an EVL, once we verify that a sequence $(u_n)_n$ satisfying (4.3) exists and that the limits p_k exist. These conditions will be addressed in Sections 5, 7 and 6.

Proof. From discussion in the previous section, in order to prove Theorem 4.1, it suffices to check conditions (R1)-(R3) for the space $(V_\alpha, \|\cdot\|_\alpha)$. The indicator function $\mathbf{1}_{\Delta_n}$ over the strip along the diagonal, $\Delta_n = \{\varphi > u_n\}$, belongs to V_α : its boundary has $(q-1)$ -dimensional Lebesgue measure $O(r_n)$ (where $r_n = e^{-u_n}$), so $|\mathbf{1}_{\Delta_n}|_\alpha = O(1)$. This together with the fact that V_α is a Banach algebra implies (R1). It is clear that all the assumptions of Lemma 3.2 are satisfied, so (R3) holds. It remains to check condition (R2). Starting by condition (B3), we see that

$$\begin{aligned} \|P_n f - P_0 f\|_{L^1} &\leq \|P_0(\mathbf{1}_{\Delta_n} f)\|_{L^1} = \|\mathbf{1}_{\Delta_n} f\|_{L^1} \\ &\leq \|f\|_{L^\infty} \cdot \|\mathbf{1}_{\Delta_n}\|_{L^1} \leq C\mu(\Delta_n) \|f\|_\alpha. \end{aligned}$$

This means that condition (B3) is satisfied, with $\pi_n = C\mu(\Delta_n)$. Condition (B2.2) follows from the fact that

$$\|P_n^k f\|_{L^1} = \|P_0^k (\mathbb{1}_{\Delta_n^c} f)\|_{L^1} = \|\mathbb{1}_{\Delta_n^c} f\|_{L^1} \leq \|f\|_{L^1}.$$

To prove condition (B2.3), we note that, using LY:

$$\begin{aligned} \|P_n f\|_\alpha &= \|P_0 (\mathbb{1}_{\Delta_n^c} f)\|_\alpha \leq \eta \|\mathbb{1}_{\Delta_n^c} f\|_\alpha + D \|\mathbb{1}_{\Delta_n^c} f\|_{L^1} \\ &\leq \eta \|\mathbb{1}_{\Delta_n^c} f\|_\alpha + (1 + D) \|f\|_{L^1}. \end{aligned}$$

To estimate $\|\mathbb{1}_{\Delta_n^c} f\|_\alpha$ we proceed as in [18, p3326] and we obtain

$$\|\mathbb{1}_{\Delta_n^c} f\|_\alpha \leq \|f\|_\alpha (1 + C_q r_n),$$

where C_q depends only on the dimension q and the parameter ϵ_0 , and $r_n = e^{-u_n}$. Condition (B2.3) then follows for all large $n \geq n_0$ by setting $r = \eta(1 + C_q r_{n_0}) < 1$ and by iterating the inequality for P_n^k . Condition (B2.1) follows from the fact that the unit ball of $(V_\alpha, \|\cdot\|_\alpha)$ is compact in the L^1 -norm. $\square$

4.3. Application to the product of piecewise expanding maps. In practice, to check the assumptions of Theorem 4.1, we will need several results from Saussol [33] that we recall below.

We say that a map $\tilde{T} : [0, 1]^q \rightarrow [0, 1]^q$ is piecewise expanding if there exists an at most countable family of disjoint open sets $U_j \subseteq [0, 1]^q$ and V_j such that $\overline{U_j} \subseteq V_j$ and maps $\tilde{T}_j : V_j \rightarrow \mathbb{R}^n$ satisfying, for some $0 < \alpha \leq 1$ and $\epsilon_0 > 0$, the following properties:

(PE1) $\tilde{T}_j|_{U_j} = \tilde{T}|_{U_j}$.

(PE2) For all j , $\tilde{T}_j \in C^1(V_j)$, $\tilde{T}_j$ is injective and $\tilde{T}_j^{-1} \in C^1(\tilde{T}_j(V_j))$. Moreover, the determinant is uniformly Hölder: for all $i, \epsilon \leq \epsilon_0, z \in \tilde{T}_j(V_j)$ and $x, y \in B_\epsilon(z) \cap \tilde{T}_j(V_j)$,

$$\left| \det D_x \tilde{T}_j^{-1} - \det D_y \tilde{T}_j^{-1} \right| \leq c \left| \det D_z \tilde{T}_j^{-1} \right| \epsilon^\alpha. \quad (4.6)$$

(PE3) $\text{Leb}([0, 1]^q \setminus \bigcup_j U_j) = 0$.

(PE4) There exists $\sigma < 1$ such that for all $u, v \in \tilde{T}(V_j)$, with $d(u, v) \leq \epsilon_0$, we have $d(T_j^{-1}u, T_j^{-1}v) \leq \sigma d(u, v)$.

(PE5) Define $G(\epsilon, \epsilon_0) := \sup_{x \in [0, 1]^q} G(x, \epsilon, \epsilon_0)$, where

$$G(x, \epsilon, \epsilon_0) := \sum_j \frac{\text{Leb}(T_j^{-1} B_\epsilon(\partial T(U_j)) \cap B_{(1-s)\epsilon_0}(x))}{\text{Leb}(B_{(1-s)\epsilon_0}(x))}.$$

The quantity $\eta(\epsilon_0) := \sigma^\alpha + 2 \sup_{\epsilon \leq \epsilon_0} \frac{G(\epsilon)}{\epsilon^\alpha} \epsilon_0^\alpha$ is such that $\sup_{\delta \leq \epsilon_0} \eta(\delta) < 1$.

In order to check condition LY, we have the following lemma.

Lemma 4.3 ([33, Lemma 4.1]). *Let P be the transfer operator associated with a map $\tilde{T}$ satisfying (PE1)-(PE5). Then, provided ϵ_0 is small enough, there exist $\eta \in (0, 1)$ and $D < \infty$ such that, for all $f \in V_\alpha$, we have that $Pf \in V_\alpha$, with*

$$\|Pf\|_\alpha \leq \eta \|f\|_\alpha + D \|f\|_{L^1([0, 1]^q, \text{Leb})}. \quad (4.7)$$

A product map $\tilde{T} : [0, 1]^q \rightarrow [0, 1]^q$ such that each factor $T_i : [0, 1] \rightarrow [0, 1]$ is piecewise $C^{1+\alpha}$ with finitely many monotonicity intervals and uniformly expanding, i.e. $\inf |T'_i| > 1$, clearly satisfies conditions (PE1)–(PE4).

Regarding condition (PE5), [33, Lemma 2.1] shows that the condition is satisfied whenever

$$\eta := \sigma^\alpha + \frac{4\sigma}{1-\sigma} Y \frac{\gamma_{q-1}}{\gamma_q} < 1,$$

provided that the dynamical partition is finite and its boundaries are contained in piecewise C^1 embedded compact codimension-one submanifolds. This applies in particular to the present product-map setting. Here, γ_q denotes the volume of the unit ball in $\mathbb{R}^q$, and Y is the maximal number of smooth boundary components of the domains of injectivity intersecting at a single point. In the present setting, we have $Y = 2q$.

Remark 4.4. In the case where all the factor maps T_i are piecewise onto, it is not necessary to take the discontinuity into consideration, and we can directly set $\eta = \sigma^\alpha$, see for example the computations in [2].

According to [33, Theorem 5.1], if the map $\tilde{T}$ satisfies conditions (PE1)–(PE5), its transfer operator is quasi-compact on V_α , and the map $\tilde{T}$ admits a spectral decomposition. In order to guarantee that there exists a unique ergodic and mixing component, and then to guarantee condition (B1), we will use the following condition (adapted from [1, Proposition 2.9]) which imposes some sort of topological exactness, and also ensures that the unique invariant density is bounded uniformly away from 0.

Proposition 4.5. *Suppose that the partition U_j associated to the map $\tilde{T}$ is finite. Assume that for any ball B of $[0, 1]^q$, there exists $n \in \mathbb{N}$ such that $\tilde{T}^n B = [0, 1]^q$ (up to Leb-measure zero), then there exists a unique absolutely continuous invariant probability (acip) $h \in V_\alpha$ for $\tilde{T}$. Furthermore, the support of h is $[0, 1]^q$ and h is bounded away from zero.*

Proof. For h an invariant density in V_α , take B to be the ball described in Lemma [33, Lemma 3.1]. Let c be the infimum of $h|_B$. Then, for Leb-almost every $x \in [0, 1]^q$,

$$h(x) = P_0^n h(x) \geq c P_0^n (\mathbf{1}_B)(x) = c \sum_j \mathbf{1}_{\tilde{T}_j^n B}(x) \left| \det D_x \left(\tilde{T}_j^n \right)^{-1} \right| \mathbf{1}_{\tilde{T}_j^n U_{i,n}}(x).$$

Here, $U_{i,n}$ denotes the partition associated to T^n . Since $\tilde{T}^n B = [0, 1]^q$, there must exist some i such that $x \in \tilde{T}_j^n B$. This means that $\left(\tilde{T}_j^n \right)^{-1}(x) \in B \cap U_{i,n}$, so $x \in \tilde{T}_j^n(U_{i,n})$. Therefore,

$$h \geq c \inf_j \inf_x \left| \det D_x \left(\tilde{T}_j^n \right)^{-1} \right| > 0, \quad (4.8)$$

which concludes the proof. $\square$

5. UNCLUSTERED SYNCHRONISATION PROFILES

Throughout this section, we assume that $q = 2$, so $\mathcal{M} = [0, 1]^2$. The product map $\bar{T}$ has smooth branches denoted by $\bar{T}_j : [0, 1]^2 \rightarrow [0, 1]^2$, and $T_{j,1}$ and $T_{j,2}$ denote the respective first and second factor maps.

In Section 7, we will study examples where the assumptions of Theorem 4.1 are verified. Here, we will address first the issue of estimating the quantities p_k given in (4.4).

We assume that our dynamical system $\bar{T} : [0, 1]^2 \rightarrow [0, 1]^2$ satisfies all the conditions of Theorem 4.1, with $\alpha = 1$. However, we recover the notation used to describe conditions (PE1)-(PE5). For example, we write that $\bar{T}_j|_{\bar{U}_j}$ is a bijection between $\bar{U}_j$ and $[0, 1]^2$. The component maps are assumed to be piecewise C^2 unidimensional expanding. This implies that each of the factor maps has a BV $([0, 1])$ invariant probability absolutely continuous with respect to Leb, which we call h_1 and h_2 . The invariant density of the product map $\bar{T}$ is

$$h : [0, 1]^2 \rightarrow [0, \infty), (x, y) \mapsto h_1(x) h_2(y).$$

The sets Δ_n can be chosen in the following way:

$$\Delta_n = \{(x, y) \in [0, 1]^2 : d(x, y) < \eta s/n\},$$

where

$$\eta = \frac{1}{2 \int_0^1 h(x, x) dx} = \frac{1}{2 \int_0^1 h_1(x) h_2(x) dx}.$$

5.1. Preparatory results. Let D be the diagonal of $[0, 1]^2$. We need this simple but useful result:

Lemma 5.1. *We have the following inclusions:*

- $\Delta_n \subseteq \{(x, y) : d((x, y), D) < \eta s/n\}$
- $\bar{T}_j|_{\bar{U}_j}^{-1}(\Delta_n) \subseteq \{(x, y) \in [0, 1]^2 : d((x, y), \bar{T}_j^{-1}D) < \eta s/n\}$, in case $\eta s/n \leq \varepsilon_0$.

Proof. To prove the first inclusion, we note that, since d is the Euclidean distance,

$$(x, y) \in \Delta_n \iff d(x, y) < \eta s/n \implies d((x, x), (x, y)) < \eta s/n.$$

Regarding the second inclusion, if $\bar{T}_j(x, y) \in \Delta_n$ then, by the first inclusion, there exists $(w, w) \in D$ such that $d(\bar{T}_j(x, y), (w, w)) < \eta s/n$. But then

$$\begin{aligned} d((x, y), \bar{T}_j^{-1}(w, w)) &= d(\bar{T}_j^{-1}\bar{T}_j(x, y), \bar{T}_j^{-1}(w, w)) \\ &\leq \sigma d(\bar{T}_j(x, y), (w, w)) < \sigma \cdot \eta s/n < \eta s/n, \end{aligned}$$

which concludes the proof. $\square$

Under the conditions described above, we can prove the following result:

Proposition 5.2. *If $\bar{T}_j|_{\bar{U}_j}^{-1}D \cap D = \emptyset$, then there exists n_0 such that for all $n \geq n_0$, the intersection $\bar{T}_j|_{\bar{U}_j}^{-1}\Delta_n \cap \Delta_n$ is empty.*

Proof. We first observe that the sets Δ_n and $\bar{T}_j|_{\bar{U}_j}^{-1}\Delta_n$ are shrinking neighborhoods of D and $\bar{T}_j|_{\bar{U}_j}^{-1}D$, respectively. Since these sets are compact and disjoint, the distance between them is strictly positive, so we can find n_0 such that for all $n \geq n_0$, the intersection $\bar{T}_j|_{\bar{U}_j}^{-1}\Delta_n \cap \Delta_n$ is empty. $\square$

5.2. Intersecting points with distinct derivatives. Before going through more results, it will be useful to study the following inequality of real numbers

$$\left| az + b \frac{z^2}{2} \right| \geq \frac{|za|}{2}. \quad (5.1)$$

If $ab = 0$, then every real number satisfies the inequality. On the other hand, if $ab > 0$, the set of solutions of the inequality is $(-\infty, -3a/b] \cup [-a/b, \infty)$. In the case of $ab < 0$, this set is $(-\infty, -a/b] \cup [-3a/b, \infty)$. Hence, the inequality is satisfied for all $z \in \mathbb{R}$, such that $|z| \leq \left| \frac{a}{b} \right|$. Another important remark is that if we replace b by b' in the inequality, where $|b'| \leq |b|$, then the set $\{z \in \mathbb{R} : |z| \leq \left| \frac{a}{b} \right|\}$ is still contained in the set of solutions of the inequality.

Recall that the factors of $\bar{T}_j : V_j \rightarrow \mathbb{R}^2$ are denoted by $T_{j,1}$ and $T_{j,2}$. Since the domains of the maps $\bar{T}_j$ are rectangles, it is not difficult to see what the domains of the maps $T_{j,1}$ and $T_{j,2}$ are intervals. We now have all the ingredients to prove a lemma, which, in loose terms, states that if there is an intersection of D with $\bar{T}^{-1}D$ at the point (x_0, x_0) , with $T'_{j,1}(x_0) \neq T'_{j,2}(x_0)$, then $\bar{T}^{-1}D$ “moves away from” D sufficiently fast.

Lemma 5.3. *Assume $\bar{T}$ is as described in the beginning of this section. Assume that $(x_0, x_0) \in \bar{U}_j$ is such that $T_{j,1}(x_0) = T_{j,2}(x_0)$, but $T'_{j,1}(x_0) \neq T'_{j,2}(x_0)$. Define*

$$\begin{aligned} \bullet \quad b_j &:= \sup_{(x,y) \in [0,1]^2} \left| \left[\left(T_{j,1}^{-1} \right)'(x) \right]^3 T''_{j,1} \left(T_{j,1}^{-1}(x) \right) - \left[\left(T_{j,2}^{-1} \right)'(y) \right]^3 T''_{j,2} \left(T_{j,2}^{-1}(y) \right) \right| \\ \bullet \quad c_j &:= \frac{\left| \left(T_{j,1}^{-1} \right)'(T_{j,1}(x_0)) - \left(T_{j,2}^{-1} \right)'(T_{j,2}(x_0)) \right|}{b_j}. \end{aligned}$$

Then, for all t such that $|t| \leq c_j$ and $(T_{j,1}(x_0) + t, T_{j,2}(x_0) + t) \in \bar{T}_j(\bar{U}_j)$, we have that

$$\left| T_{j,1}^{-1}(T_{j,1}(x_0) + t) - T_{j,2}^{-1}(T_{j,2}(x_0) + t) \right| \geq \frac{|t|}{2} \left| \left(T_{j,1}^{-1} \right)'(T_{j,1}(x_0)) - \left(T_{j,2}^{-1} \right)'(T_{j,2}(x_0)) \right|.$$

Proof. By Taylor expansion, we have

$$\begin{aligned} \left| T_{j,1}^{-1}(T_{j,1}(x_0) + t) - T_{j,2}^{-1}(T_{j,2}(x_0) + t) \right| &= \\ \left| \left[\left(T_{j,1}^{-1} \right)'(T_{j,1}(x_0)) - \left(T_{j,2}^{-1} \right)'(T_{j,2}(x_0)) \right] t \right. & \\ \left. + \left[\left(T_{j,1}^{-1} \right)''(T_{j,1}(x_0) + r_1) - \left(T_{j,2}^{-1} \right)''(T_{j,2}(x_0) + r_2) \right] \frac{t^2}{2} \right|. & \end{aligned}$$

The result follows by applying the observation above about the inequality (5.1), with

$$a = \left(T_{j,1}^{-1} \right)'(T_{j,1}(x_0)) - \left(T_{j,2}^{-1} \right)'(T_{j,2}(x_0))$$

and

$$b = \sup_{(x,y) \in [0,1]^2} \left| \left[\left(T_{j,1}^{-1} \right)'(x) \right]^3 T''_{j,1} \left(T_{j,1}^{-1}(x) \right) - \left[\left(T_{j,2}^{-1} \right)'(y) \right]^3 T''_{j,2} \left(T_{j,2}^{-1}(y) \right) \right|.$$

The upper bound b is obtained by differentiating twice the expression $f(f^{-1}(x)) = x$. $\square$

As a consequence we obtain the following corollary.

Corollary 5.4. *Let*

$$0 < \varepsilon < \frac{\left| (T_{j,1}^{-1})'(T_{j,1}(x_0)) - (T_{j,2}^{-1})'(T_{j,2}(x_0)) \right|^2}{2b_j} \quad \text{and} \quad \delta = \frac{2\varepsilon}{\left| (T_{j,1}^{-1})'(T_{j,1}(x_0)) - (T_{j,2}^{-1})'(T_{j,2}(x_0)) \right|}.$$

Then, for all $t \in (\delta, c_j) \cup (-c_j, -\delta)$, we have

$$\left| T_{j,1}^{-1}(T_{j,1}(x_0) + t) - T_{j,2}^{-1}(T_{j,2}(x_0) + t) \right| \geq \varepsilon.$$

5.3. Isolated intersecting points have negligible contribution for the computation of θ . Now we prove that under the conditions of Lemma 5.3, (x_0, x_0) is an isolated point of $D \cap \bar{T}^{-1}D$. Moreover, its contribution for the computation of the extremal index is negligible. From now on, by $\bar{T}_j^{-1}$, we mean $\bar{T}_j^{-1}|_{\bar{U}_j}$.

Proposition 5.5. *Suppose that $\bar{T}$ satisfies the properties described in the beginning of this section and that, for some j , $(x_0, x_0) \in \bar{U}_j \cap D \cap \bar{T}_j^{-1}(D)$ is a point for which $T'_{j,1}(x_0) \neq T'_{j,2}(x_0)$. Then there exists a neighbourhood V of (x_0, x_0) such that*

$$V \cap D \cap \bar{T}_j^{-1}D = \{(x_0, x_0)\} \quad \text{and} \quad V \cap D \cap \bar{T}^{-1}D \subseteq \{(x_0, x_0)\}.$$

Furthermore,

$$\lim_{n \rightarrow \infty} \frac{\mu(\Delta_n \cap \bar{T}^{-1}\Delta_n \cap V)}{\mu(\Delta_n)} = 0.$$

Remark 5.6. The inclusion $V \cap D \cap \bar{T}^{-1}D \subseteq \{(x_0, x_0)\}$ may not be an equality because (x_0, x_0) may belong to ∂U_j , where the map can be defined arbitrarily.

Proof. First, suppose that $(x_0, x_0) \in U_j$. We claim that $V = T_j^{-1}(B_{c_j}(T_{j,1}x_0, T_{j,2}x_0))$ is the desired neighborhood. Note that this neighborhood is not necessarily an open set of $[0, 1]^2$, but it is on the induced topology of $\bar{U}_j$. To prove this claim, we choose $(x, y) \in T_j^{-1}D \cap V \setminus \{(x_0, x_0)\}$. We know that there exists t , where $0 < |t| < c_j$, such that $(x, y) = (T_{j,1}^{-1}(T_{j,1}x_0 + t), T_{j,2}^{-1}(T_{j,2}x_0 + t))$. Lemma 5.3 tells us $(x, y) \notin D$. If $(x_0, x_0) \in \partial U_j$, one should take all i for which $(x_0, x_0) \in D \cap T_j^{-1}D$ and the neighborhood V_j of that point as above. For the i such that $(x_0, x_0) \notin T_j^{-1}D$, we can take the ball B_{r_j} centered at (x_0, x_0) of radius $r_j := d((x_0, x_0), T_j^{-1}D)/2$. The union of all of these gives a neighborhood of (x_0, x_0) with the desired conditions. Of course, we can ensure these neighborhoods are open by taking a ball contained in V . One important thing to mention is that

$$V \subseteq \bigcup_{j: (x_0, x_0) \in \bar{U}_j} \bar{U}_j.$$

For each j such that $(x_0, x_0) \in \bar{U}_j$, define the following quantities:

$$\Gamma_j = \frac{\left| (T_{j,1}^{-1})'(T_{j,1}x_0) - (T_{j,2}^{-1})'(T_{j,2}x_0) \right|^2}{2b_j} \tag{5.2}$$

$$\Lambda_j = \frac{\epsilon_0 \left| (T_{j,1}^{-1})'(T_{j,1}x_0) - (T_{j,2}^{-1})'(T_{j,2}x_0) \right|}{2\sqrt{2}} \tag{5.3}$$

we take ε_j as follows:

$$\varepsilon_j < \begin{cases} \min \{3\epsilon_0, 3r_j\} & (x_0, x_0) \notin T_j^{-1}D \\ \min \left\{ c_j, \Gamma_j, \frac{3d_j}{2}, \Lambda_j \right\}, & (x_0, x_0) \in T_j^{-1}D \end{cases} \quad (5.4)$$

where $d_j := d\left(T_j^{-1}D, D \cap T_j^{-1}\left(\overline{B_{c_j}(T_{j,1}x_0, T_{j,2}x_0)} \setminus B_{c_j/2}(T_{j,1}x_0, T_{j,2}x_0)\right)\right)$, which is greater than 0, since both sets are compact and do not intersect each other. Let $\varepsilon = \min_{i: (x_0, x_0) \in \overline{U_j}} \varepsilon_j$.

Suppose that $(x, y) \in U_j$ where i is such that $(x_0, x_0) \in T_j^{-1}D$. Additionally, suppose $(x, y) \in \Delta_n \cap T_j^{-1}\Delta_n \cap T_j^{-1}\left[B_{c_j/2}(T_{j,1}x_0, T_{j,2}x_0)\right]$ for sufficiently large n (i.e., $1/n \leq \varepsilon/(3\eta s)$). We know that $(T_{j,1}x, T_{j,2}y) \in \Delta_n$, which implies, by Lemma 5.1, that $d((T_{j,1}x, T_{j,2}y), D) < s\varepsilon/3$. Let $(z, z) \in D$ be such that

$$d((T_{j,1}x, T_{j,2}y), (z, z)) < s\varepsilon/3.$$

Then,

$$\begin{aligned} d((T_{j,1}x_0, T_{j,2}x_0), (z, z)) &\leq d((T_{j,1}x_0, T_{j,2}x_0), (T_{j,1}x, T_{j,2}y)) + d((T_{j,1}x, T_{j,2}y), (z, z)) \\ &\leq \frac{c_j}{2} + \frac{s\varepsilon}{3} \leq \frac{c_j}{2} + \frac{\varepsilon}{3} < c_j. \end{aligned}$$

This means that $(z, z) \in B_{c_j}(T_{j,1}x_0, T_{j,2}x_0)$. Let $(u, v) = \bar{T}_j^{-1}(z, z)$. Since

$$(u, v) \in \bar{T}_j^{-1}B_{c_j}(T_{j,1}x_0, T_{j,2}x_0),$$

there exists $|t| < c_j$ such that

$$(u, v) = \left(T_{j,1}^{-1}(T_{j,1}x_0 + t), T_{j,2}^{-1}(T_{j,2}x_0 + t)\right).$$

Note that

$$d(u, v) \leq d(u, x) + d(x, y) + d(y, v) < s/n \leq \varepsilon.$$

Since $\varepsilon < \frac{\left|(T_{j,1}^{-1})'(T_{j,1}x_0) - (T_{j,2}^{-1})'(T_{j,2}x_0)\right|^2}{2b_j}$, Corollary 5.4 tells us that

$$|t| < \delta = \frac{2\varepsilon}{\left|\left(T_{j,1}^{-1}\right)'(T_{j,1}x_0) - \left(T_{j,2}^{-1}\right)'(T_{j,2}x_0)\right|}.$$

Therefore,

$$d((z, z), (T_{j,1}x_0, T_{j,2}x_0)) < \sqrt{2} \cdot \delta.$$

Since by the choice of ε , we have $\sqrt{2} \cdot \delta < \epsilon_0$, then

$$\begin{aligned} d((u, v), (x_0, x_0)) &= d\left(\bar{T}_j^{-1}(z, z), T_j^{-1}(T_{j,1}x_0, T_{j,2}x_0)\right) \\ &\leq \sigma d((z, z), (T_{j,1}x_0, T_{j,2}x_0)) < \sqrt{2} \cdot \delta. \end{aligned}$$

Finally,

$$\begin{aligned} d((x, y), (x_0, x_0)) &\leq d((x, y), (u, v)) + d((u, v), (x_0, x_0)) < \\ &\varepsilon \left(\frac{1}{3} + \frac{2\sqrt{2}}{\left|\left(T_{j,1}^{-1}\right)'(T_{j,1}x_0) - \left(T_{j,2}^{-1}\right)'(T_{j,2}x_0)\right|} \right). \end{aligned}$$

On the other hand, if $(x, y) \in T_j^{-1} \left[B_{c_j}(T_{j,1}x_0, T_{j,2}x_0) \setminus B_{c_j/2}(T_{j,1}x_0, T_{j,2}x_0) \right]$, then

$$d((x, y), D) \geq \varepsilon/3 \text{ or } d((x, y), T_j^{-1}D) \geq \varepsilon/3.$$

Otherwise, by the triangle inequality, we would get $d_j \leq \frac{2\varepsilon}{3}$, contradicting the choice of ε . So, by Lemma 5.1, we get $(x, y) \notin \Delta_n \cap \bar{T}_j^{-1}\Delta_n$.

In case $(x_0, x_0) \notin \bar{T}_j^{-1}D$, we have that

$$\bar{T}_j^{-1}\Delta_n \subseteq \left\{ (x, y) \in \bar{U}_j : d((x, y), \bar{T}_j^{-1}D) < \varepsilon/3 \right\}.$$

Note that

$$2r_j = d((x_0, x_0), \bar{T}_j^{-1}D) \leq d((x_0, x_0), (x, y)) + d((x, y), \bar{T}_j^{-1}D),$$

which means that if $(x, y) \in V \cap U_j$, then

$$d((x, y), \bar{T}_j^{-1}D) \geq r_j > \varepsilon/3,$$

which implies that $(x, y) \notin \bar{T}_j^{-1}\Delta_n$.

In conclusion,

$$\mu(\Delta_n \cap \bar{T}^{-1}\Delta_n \cap V) \leq \sum_{i: (x_0, x_0) \in \bar{U}_j} \mu(\Delta_n \cap \bar{T}_j^{-1}\Delta_n \cap V) = \mathcal{O}(n^{-2}),$$

since by the positivity assumption and the fact that $h \in V_\alpha \subseteq L^\infty$,

$$\mu(B_{1/n}(x_0, x_0)) = \mathcal{O}(n^{-2}).$$

The conclusion follows by the fact that $\mu(\Delta_n) = \mathcal{O}(n^{-1})$. $\square$

5.4. Corollaries and sufficient conditions for $\theta = 1$. Now, we point out some corollaries of Proposition 5.5. An important thing to mention is that if $\bar{T}$ satisfies the properties described in the beginning of this section, so do the iterates of $\bar{T}$. This means that we can also apply Proposition 5.5 to the iterates of $\bar{T}$. With this in mind, we have the following:

Corollary 5.7. *Suppose that for all $\kappa \in \mathbb{N}$, the number of points of $\bigcup_j \bar{T}_j^{-\kappa}D \cap D$ is finite. If for all these points the derivatives of the components are different, then the extremal index is 1.*

Proof. For each $(x_k, x_k) \in \bigcup_j \bar{T}_j^{-1}D \cap D$, let V_k be the neighborhood as in Proposition 5.5, which we assume, without loss of generality, to be an open set of $[0, 1]^2$. Let $W = (\bigcup_k V_k)^c$. We note that $\bigcup_j \bar{T}_j^{-1}D \cap W$ is a compact set whose intersection with D is empty. So we can take n_0 large enough such that

$$\bigcup_j \bar{T}_j^{-1}\Delta_n \cap \Delta_n \cap W = \emptyset \quad \text{for all } n \geq n_0.$$

Note that, up to Lebesgue measure zero, and consequently μ -measure zero,

$$\bigcup_j \bar{T}_j^{-1}\Delta_n = \bar{T}^{-1}\Delta_n.$$

This allows us to make the following estimation

$$\mu(\bar{T}^{-1}\Delta_n \cap \Delta_n) \leq \sum_k \mu(\bar{T}^{-1}\Delta_n \cap \Delta_n \cap V_k) + \mu(\bar{T}^{-1}\Delta_n \cap \Delta_n \cap W).$$

Dividing by $\mu(\Delta_n)$ and using Proposition 5.5, we conclude that $p_0 = 0$, where p_κ is as defined in (3.5). For $\kappa \geq 0$, we can apply the same reasoning. Since

$$\mu(\Delta_n \cap \bar{T}^{-1}\Delta_n^c \cap \dots \cap \bar{T}^{-\kappa}\Delta_n^c \cap \bar{T}^{-(\kappa+1)}\Delta_n) \leq \mu(\Delta_n \cap \bar{T}^{-(\kappa+1)}\Delta_n),$$

it follows that $p_\kappa = 0$. $\square$

Corollary 5.8. *Let $\ell \geq 1$. Assume that there exists $k_\ell > 0$ such that for all intersection points $(x_0, x_0) \in D \cap \bigcup_j \bar{T}_j^{-\ell} D$,*

$$\left| \left(T_{j,1}^{-\ell} \right)' (T_{j,1}^\ell x_0) - \left(T_{j,2}^{-\ell} \right)' (T_{j,2}^\ell x_0) \right| > k_\ell.$$

Then, the number of points of $D \cap \bigcup_j \bar{T}_j^{-\ell} D$ is finite, which implies that $p_{\ell-1} = 0$.

Proof. It is enough to prove that, for each i , $D \cap \bar{T}_j^{-1} D$ is finite. By the proof of the proposition, if $(x_0, x_0) \in D \cap \bar{T}_j^{-1} D$, then it is the unique point of that set when intersected with $\bar{T}_j^{-1}(B_{c_j}(T_{j,1}x_0, T_{j,2}x_0))$. But $c_j > k_1/b_j$, so (x_0, x_0) is the unique point in

$$D \cap \bar{T}_j^{-1} D \cap \bar{T}_j^{-1} \left(B_{k_1/b_j}(T_{j,1}x_0, T_{j,2}x_0) \right).$$

What remains to prove is that, for all $\varepsilon > 0$, there is $l_\varepsilon > 0$ such that for all $(x, x) \in \bar{U}_j \cap D$, the curve $\bar{T}_j^{-1}[B_\varepsilon(x, x) \cap D]$ has a length greater than l_ε . This means that if (x_0, x_0) and (x_1, x_1) are different points of $\bar{T}_j^{-1} D$, then $(x_1, x_1) \notin \bar{T}_j^{-1} \left(B_{k_1/b_j}(T_{j,1}x_0, T_{j,2}x_0) \right)$, which implies, by the construction of the proof, that the path in $\bar{T}_j^{-1} D$ whose end-points are (x_0, x_0) and (x_1, x_1) has length greater than $l_{k_1/b_j}/2$. This implies that there is only a finite number of such points. To get the lower bound for the length of the curve, we note that

$$l\left(\bar{T}_j^{-1}[B_\varepsilon(x, x) \cap D]\right) = \int_{-\varepsilon}^{\varepsilon} \sqrt{\left[\left(T_{j,1}^{-1} \right)'(x+t) \right]^2 + \left[\left(T_{j,2}^{-1} \right)'(x+t) \right]^2} dt.$$

The domain of integration might not be exactly $(-\varepsilon, \varepsilon)$ as $B_\varepsilon(x, x) \cap D$ might not be contained in $\bar{U}_j$. But it will always be an interval of length at least ε . Suppose that the domain of integration is contained in $\bar{U}_j$. Then, using the fact that $\bar{T}_j$ is C^1 and invertible in an open set containing $\bar{U}_j$, we conclude, by the formula of the derivative of the inverse function, that $\left(T_{j,1}^{-1} \right)'$ and $\left(T_{j,2}^{-1} \right)'$ are both bounded away from zero, allowing us to obtain a lower bound for $l\left(\bar{T}_j^{-1}[B_\varepsilon(x, x) \cap D]\right)$. $\square$

6. THE EXTREMAL INDEX AS A MEASURE OF DYNAMICAL COMPATIBILITY

In this section, we show how the Extremal Index distinguishes between compatible and incompatible factor dynamics, quantifying the degree of compatibility.

We will consider 2-dimensional linear dynamics, which will simplify the computation of the coefficients p_k given in (3.5), which are crucial to compute the Extremal Index. These systems

clearly fit the framework of Theorem 4.1, as they can be seen as particular cases of the class of maps studied in Section 4.3, for example.

We will start by verifying that if the linear maps have different coefficients then the Extremal Index is 1, which could be obtained easily from the results in Section 5. However, we perform a direct analysis in order to provide further insight about what is behind this interpretation of the role of the Extremal Index and at the same time take the opportunity to set the notation and establish a benchmark against which the cases of full and partial compatibility can be compared.

Full compatibility is obtained by considering the same linear coefficients for the two factor maps, which will give rise to an Extremal Index less than 1, where we recover the formula obtained in previous works, [10]. Then, we will introduce an example of different linear factor maps, which coincide in some part of phase space, creating a relevant compatibility detected by an Extremal Index less than 1, which will turn out to be a weighted average of the two previous cases.

6.1. Direct product of different linear maps. Consider the map

$$\begin{aligned} T: [0, 1]^2 &\longrightarrow [0, 1]^2 \\ (x, y) &\longmapsto (\mathbf{a}x \bmod 1, \mathbf{b}y \bmod 1) \end{aligned} \quad (6.1)$$

where $\mathbf{a}, \mathbf{b} \in \mathbb{Z} \setminus \{-1, 0, 1\}$. The invariant measure μ is Lebesgue measure. In this case, the observable function giving rise to the stochastic process, can be written in the simplified form:

$$\begin{aligned} \varphi: [0, 1]^2 &\longrightarrow \mathbb{R} \\ (x, y) &\longmapsto -\log d(x, y) \end{aligned} \quad (6.2)$$

For each $\tau > 0$, define

$$u_n(\tau) = \log \frac{2n}{\tau}. \quad (6.3)$$

It is clear that the set $\Delta_n = \{\varphi > u_n\} = \left\{ (x, y) \in [0, 1]^2 : d(x, y) < \frac{\tau}{2n} \right\}$. We denote the quantity $\frac{\tau}{2n}$ by r_n . It is also clear, since μ is the Lebesgue measure, that $\mu(\Delta_n) \sim \frac{\tau}{n}$. The map T can be partitioned into $|\mathbf{a}\mathbf{b}|$ open sets, where T restricted to these parts is a diffeomorphism. Instead of denoting these elements by U_i as in the previous chapter, we will denote them by I_{kl} , where $k \in \{0, \dots, \mathbf{a} - 1\}$ or $k \in \{-1, \dots, \mathbf{a}\}$ and $l \in \{0, \dots, \mathbf{b} - 1\}$ or $l \in \{-1, \dots, \mathbf{b}\}$, depending on whether $\mathbf{a}$ and $\mathbf{b}$ are negative or positive. Similarly, we denote the elements of the analogous partition T^j by I_{kl}^j , where $0 \leq k \leq \mathbf{a}^j - 1$ or $-1 \geq k \geq \mathbf{a}^j$, and $0 \leq l \leq \mathbf{b}^j - 1$ or $-1 \geq l \geq \mathbf{b}^j$, depending on whether $\mathbf{a}^j$ and $\mathbf{b}^j$ are positive or negative (the sign that matters is the one of $\mathbf{a}^j$, not $\mathbf{a}$). Explicitly,

$$I_{kl}^j = \left\{ (x, y) \in [0, 1]^2 : x \in I_k^j, y \in J_l^j \right\}, \quad (6.4)$$

where $I_k^j = \left(\frac{k}{\mathbf{a}^j}, \frac{k+1}{\mathbf{a}^j} \right)$ or $I_k^j = \left(\frac{k+1}{\mathbf{a}^j}, \frac{k}{\mathbf{a}^j} \right)$, depending on whether $\mathbf{a}^j$ is positive or negative. The set J_l^j is defined similarly, but replacing $\mathbf{a}$ by $\mathbf{b}$. We define the maps

$$T_{kl}^j: \mathbb{R}^2 \rightarrow \mathbb{R}^2, (x, y) \mapsto (\mathbf{a}^j x - k, \mathbf{b}^j y - l). \quad (6.5)$$

Clearly, $T^j|_{I_{kl}^j} = T_{kl}^j|_{I_{kl}^j}$. We denote the diagonal of $[0, 1]^2$ by D .

In this section, we verify that when $|\mathbf{a}| \neq |\mathbf{b}|$, then Extremal Index is $\theta = 1$. This can be explained by the fact that, for a fixed $j \in \mathbb{N}$, there is only a finite number of points on the diagonal that return to the diagonal after j iterations. Since the derivatives of the component maps are different, then the area of a small neighbourhood of those points that return to a neighbourhood of the diagonal is negligible for the computation of the Extremal Index.

The proof will be based on checking that the following condition holds:

$$\lim_{n \rightarrow \infty} n\mu(\Delta_n \cap T^{-j}(\Delta_n)) = 0, \quad \text{for all } j \in \mathbb{N}. \quad (6.6)$$

Note that validity of (6.6) implies that $p_k = 0$, for all $k \in \mathbb{N}_0$ and therefore $\theta = 1 - \sum_{k \geq 0} p_k = 1$.

We start by noting that $T^{-j}(D) \cap I_{kl}^j$ is the line

$$\left(y - \frac{l}{\mathbf{b}^j}\right) = \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k}{\mathbf{a}^j}\right)$$

intersected with I_{kl}^j . The set $T^{-j}(\Delta_n) \cap I_{kl}^j$ is the space between the lines

$$y - \frac{l + (\frac{t}{2n})}{\mathbf{b}^j} = \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k}{\mathbf{a}^j}\right); \quad y - \frac{l}{\mathbf{b}^j} = \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k + (\frac{t}{2n})}{\mathbf{a}^j}\right).$$

So, $T^{-j}(\Delta_n) \cap I_{kl}^j$ is the set of points $(x, y) \in I_{kl}^j$ such that

$$\left| \left(y - \frac{l}{\mathbf{b}^j}\right) - \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k}{\mathbf{a}^j}\right) \right| < \frac{r_n}{|\mathbf{b}|^j}.$$

We note that if $D \cap T^{-j}(D) \cap I_{kl}^j \neq \emptyset$, then it contains only one point, since it corresponds to the intersection of two linear maps with different derivatives. To prove condition (6.6), we need an upper bound for the measure of the sets $\Delta_n \cap T^{-j}\Delta_n$. To get it, we analyse these sets restricted to I_{kl}^j .

Lemma 6.1. *Suppose that $T : [0, 1]^2 \rightarrow [0, 1]^2$ is a map and that there exists an open set A such that $T^j|_A$ is a diffeomorphism between A and $\text{int}([0, 1]^2)$ of the form $T^j|_A(x, y) = (\mathbf{a}^j x - q, \mathbf{b}^j y - p)$, with $|\mathbf{a}| \neq |\mathbf{b}|$. Suppose also that $D \cap \overline{T^{-j}(D)} \cap A = (x_0, x_0)$. Then, the set $\Delta_n \cap T^{-j}(\Delta_n) \cap A$ is a subset of*

$$\left\{ (x, y) \in A : d(x, x_0) \leq \frac{r_n(1 + |\mathbf{b}|^j)}{||\mathbf{b}|^j - |\mathbf{a}|^j|}, d(y, x_0) \leq \frac{r_n(1 + |\mathbf{a}|^j)}{||\mathbf{b}|^j - |\mathbf{a}|^j|} \right\},$$

which is non-empty.

Proof. Suppose that $(x, y) \in \Delta_n \cap T^{-j}(\Delta_n) \cap A$ and that $(x_0, x_0) \in \overline{T^{-j}D \cap A}$. We note that $T^j|_A$ can be extended to $\mathbb{R}^2$, so we will assume that $T^j(x_0, x_0) = (T_1^j x_0, T_2^j x_0)$ and $T_1^j x_0 = T_2^j x_0$, where

$$T_1^j(x) = \mathbf{a}^j x - q \quad T_2^j(x) = \mathbf{b}^j x - p.$$

We want to estimate $d(x, x_0)$ and $d(y, x_0)$. Assume first that $|\mathbf{b}|^j > |\mathbf{a}|^j$. Then,

$$\begin{aligned} d(y, x_0) &\leq \frac{d(T_2^j y, T_2^j x_0)}{|\mathbf{b}|^j} \leq \frac{d(T_1^j x, T_2^j y) + d(T_1^j x, T_1^j x_0)}{|\mathbf{b}|^j} \leq \frac{r_n + |\mathbf{a}|^j d(x, x_0)}{|\mathbf{b}|^j} \\ &\leq \frac{r_n + |\mathbf{a}|^j (d(y, x) + d(y, x_0))}{|\mathbf{b}|^j} \end{aligned}$$

Therefore,

$$d(y, x_0) \leq \frac{r_n (1 + |\mathbf{a}|^j)}{|\mathbf{b}|^j - |\mathbf{a}|^j}.$$

By a simple triangle inequality, we conclude that

$$d(x, x_0) \leq d(y, x) + d(y, x_0) \leq \frac{r_n (1 + |\mathbf{b}|^j)}{|\mathbf{b}|^j - |\mathbf{a}|^j}.$$

The proof of the case where $|\mathbf{a}|^j > |\mathbf{b}|^j$ is similar. $\square$

Define the quantity

$$\beta_n = \max \left\{ m \in \mathbb{N} : \forall 1 \leq j \leq m : T^{-j} \Delta_n \cap \Delta_n \cap I_{kl}^j \neq \emptyset \iff D \cap \overline{T^{-j} D \cap I_{kl}^j} \neq \emptyset, \right. \\ \left. \forall k, l \text{ where } I_{kl}^j \text{ is defined.} \right\} \quad (6.7)$$

Lemma 6.2. $\lim_{n \rightarrow \infty} \beta_n = +\infty$.

Proof. Note that β_n is an increasing sequence of integers. So, the only way its limit is not infinity is if the sequence is constant after a certain n . Assume that this is indeed the case. This means that there exists $j, k, l \in \mathbb{N}$ such that, for all $n \in \mathbb{N}$, $T^{-j} \Delta_n \cap \Delta_n \cap I_{kl}^j \neq \emptyset$, but $D \cap \overline{T^{-j} D \cap I_{kl}^j} = \emptyset$. Both D and $\overline{T^{-j} D \cap I_{kl}^j}$ are compact, which means that if the intersection of these sets is empty, then $d(D, \overline{T^{-j} D \cap I_{kl}^j}) > 0$. Take n such $\frac{\tau}{n} < d(D, \overline{T^{-j} D \cap I_{kl}^j})$. Then, if $(x, y) \in T^{-j} \Delta_n \cap \Delta_n \cap I_{kl}^j$, we have that

$$d(D, \overline{T^{-j} D \cap I_{kl}^j}) \leq d(D, (x, y)) + d(\overline{T^{-j} D \cap I_{kl}^j}, (x, y)) < 2 \frac{\tau}{2n} = \frac{\tau}{n},$$

which yields a contradiction. To justify the last inequality, note that:

$$(x, y) \in \Delta_n \implies |x - y| < \frac{\tau}{2n} \implies d((x, y), D) \leq \frac{\tau}{2n} = r_n,$$

and also that

$$(x, y) \in T^{-j} \Delta_n \cap I_{kl}^j \implies \left| \left(y - \frac{l}{\mathbf{b}^j} \right) - \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k}{\mathbf{a}^j} \right) \right| < \frac{r_n}{|\mathbf{b}|^j}.$$

Since $(x, y) \in I_{kl}^j$, then $\left(x, \frac{\mathbf{a}^j}{\mathbf{b}^j} \left(x - \frac{k}{\mathbf{a}^j} \right) + \frac{l}{\mathbf{b}^j} \right) \in I_{kl}^j \cap T^{-j}(D)$. This means that

$$d(T^{-j} D \cap I_{kl}^j, (x, y)) \leq \frac{r_n}{|\mathbf{b}|^j} < r_n.$$

The fact that $d(T^{-j} D \cap I_{kl}^j, (x, y)) = d(\overline{T^{-j} D \cap I_{kl}^j}, (x, y))$ concludes the proof. $\square$

We are now ready to establish the following.

Proposition 6.3. *Condition (6.6) is satisfied.*

Proof. The number of points of $\bigcup_{kl} \overline{D \cap T^{-j} D \cap I_{kl}^j}$ is bounded by $|\mathbf{a}^j - \mathbf{b}^j| + 1$, since $(x, x) \in \overline{D \cap T^{-j} D \cap I_{kl}^j} \implies \mathbf{a}^j x - \mathbf{b}^j x \in \mathbb{Z}$. Since the sets I_{kl}^j are disjoint and their union has full measure, it follows that

$$n\mu(\Delta_n \cap T^{-j} \Delta_n) = n \sum_{k,l} \mu(\Delta_n \cap T^{-j} \Delta_n \cap I_{kl}^j).$$

Noting that each point of $[0, 1]^2$ is, at most, in four of the $\overline{I_{kl}^j}$, and assuming that n is sufficiently large so that $\beta_n > j$ (which we can assume by Lemma 6.2), then we can use Lemma 6.1 to obtain

$$n\mu(\Delta_n \cap T^{-j} \Delta_n) \leq 4n \left((|\mathbf{a}^j - \mathbf{b}^j| + 1) \left(\frac{(1 + |\mathbf{a}|^j)(1 + |\mathbf{b}|^j)}{(|\mathbf{a}|^j - |\mathbf{b}|^j)^2} \right) r_n^2 \right).$$

□

6.2. Direct product of identical linear maps. In this section we consider T to be given as in (6.1), but this time we assume that $\mathbf{b} = \mathbf{a}$. Let the observable φ be as before, (6.2).

We define the quantity β_n as in (6.7). It goes to infinity as n goes to infinity, by Lemma 6.2. It is easy to see that condition (6.6) does not hold anymore, but we introduce a new version which implies that $p_k = 0$ for all $k \in \mathbb{N}$ and then we will only be left with computing p_0 in order to obtain the Extremal Index.

Let $\Delta_n^{(1)} = \Delta_n \cap T^{-1}(\Delta_n^c)$. We will show that

$$\lim_{n \rightarrow \infty} n\mu(\Delta_n^{(1)} \cap T^{-j}(\Delta_n)) = 0, \quad \text{for all } j \geq 2. \quad (6.8)$$

Proposition 6.4. *Condition (6.8) is satisfied.*

Proof. The first thing to notice is that

$$\Delta_n^{(1)} = \left\{ (x, y) \in [0, 1]^2 : \frac{r_n}{|\mathbf{a}|} < |x - y| < r_n \right\} \cup B,$$

where $m(B) = \mathcal{O}(\frac{1}{n^2})$. Note that for $j < \beta_n$ the set $I_{kl}^j \cap T^{-j}(\Delta_n) \cap \Delta_n^{(1)}$ is, in fact, empty. To see this, observe that

$$(x, y) \in T^{-j}(\Delta_n) \cap I_{kl}^j \implies \left| \left(y - \frac{l}{\mathbf{a}^j} \right) - \left(x - \frac{k}{\mathbf{a}^j} \right) \right| < \frac{r_n}{|\mathbf{a}|^j}.$$

By the choice of β_n , the intersection $I_{kl}^j \cap T^{-j}(\Delta_n) \cap \Delta_n^{(1)}$ can only be non-empty if $k = l$. But in that case,

$$(x, y) \in \Delta_n^{(1)} \cap T^{-j}(\Delta_n) \cap I_{kk}^j \implies \frac{r_n}{|\mathbf{a}|} < |x - y| < \frac{r_n}{|\mathbf{a}|^j},$$

and since $|\mathbf{a}| > 1$, $\Delta_n^{(1)} \cap T^{-j}(\Delta_n) \cap I_{kk}^j = \emptyset$. Hence, the conclusion follows trivially. □

We are now able to compute the Extremal Index.

Proposition 6.5. *The extremal index is*

$$\theta = \frac{|\mathbf{a}| - 1}{|\mathbf{a}|}.$$

Proof. Start by noting that validity of (6.8) implies that $p_k = 0$ for all $k \in \mathbb{N}$. So we are left to compute

$$p_0 = \lim_{n \rightarrow \infty} 1 - \frac{\mu(\Delta_n^{(1)})}{\mu(\Delta_n)} = 1 - \lim_{n \rightarrow \infty} \frac{\frac{(|\mathbf{a}|-1)}{|\mathbf{a}|} r_n + \mathcal{O}\left(\frac{1}{n^2}\right)}{r_n} = 1 - \frac{|\mathbf{a}| - 1}{|\mathbf{a}|}.$$

□

6.3. Direct product of partially matching linear maps. We consider now linear factor maps that coincide only in a given part of the phase space. Namely, we consider

$$\begin{aligned} T: [0, 1]^2 &\longrightarrow [0, 1]^2 \\ (x, y) &\longmapsto (T_1(x), T_2(y)) \end{aligned}$$

where

$$T_1(x) = \mathbf{a}x \mod 1 \quad \text{and} \quad T_2(y) = \begin{cases} \mathbf{a}y \mod 1, & y \leq 1/|\mathbf{a}| \\ \mathbf{a}by \mod 1, & y \geq 1/|\mathbf{a}|. \end{cases}$$

The invariant measure μ is still the Lebesgue measure and Theorem 4.1 is still valid. For convenience, we label the partition associated with T differently. Let $\{J_k\}_{k=0}^{(|\mathbf{a}|-1)|\mathbf{b}|}$ be the partition associated with T_2 . Denote by $C_{i_0, \dots, i_{j-1}}$ the j -cylinders associated with T_2 . Note that $y \in C_{i_0, \dots, i_{j-1}}$ means that $y \in J_{i_0}$, $T_2(y) \in J_{i_1}$, ..., $T_2^{j-1}(y) \in J_{i_{j-1}}$. We denote the elements of the partition associated with T_1 by $\{I_k^j\}_{k=0}^{|\mathbf{a}|^j-1}$. As in the previous case, we will prove that condition (6.8) holds.

Proposition 6.6. *Condition (6.8) is satisfied and the Extremal Index is*

$$\theta = \frac{|\mathbf{a}| - 1}{|\mathbf{a}|^2} + \frac{|\mathbf{a}| - 1}{|\mathbf{a}|}.$$

Proof. Let β_n be defined as in (6.7). The map restricted to the open set $I_0^j \times C_{00\dots 0}$ is equal to the one studied previously in this section. So, if $j < \beta_n$, it was shown in the proof of Proposition 6.4 that

$$\Delta_n^{(1)} \cap T^{-j} \Delta_n^{(1)} \cap I_0^j \times C_{00\dots 0} = \emptyset. \quad (6.9)$$

In the remaining regions, the maps are different. In fact, in those regions, the map T_2 assumes the form $\mathbf{a}^j \mathbf{b}^m y \mod 1$, for some $m \in \{1, \dots, j\}$. To get an upper bound for the number of points of the diagonal which are mapped into the diagonal, we sum

$$\sum_{m=1}^j |\mathbf{a}|^j (|\mathbf{b}|^m - 1) = \frac{|\mathbf{b}|}{|\mathbf{b}| - 1} |\mathbf{a}|^j (|\mathbf{b}|^j - 1) - j |\mathbf{a}|^j,$$

where each term of the sum represents the number of points of the diagonal mapped into the diagonal, when we consider the map $(\mathbf{a}^j x \mod 1, \mathbf{b}^m \mathbf{a}^j y \mod 1)$, after removing the point $(0, 0)$, since it is always in the region where both maps are equal.

Using Lemma 6.2, we may assume that n is sufficiently large so that $j \leq \beta_n$ and then, by Lemma 6.1, we have

$$\mu\left(\Delta_n^{(1)} \cap T^{-j}\Delta_n\right) \leq \left[\frac{|\mathbf{b}|}{|\mathbf{b}|-1} |\mathbf{a}|^j (|\mathbf{b}|^j - 1) - j |\mathbf{a}|^j\right] \cdot \left(\frac{r_n^2 (1 + |\mathbf{a}\mathbf{b}|^j) \cdot (1 + |\mathbf{a}|^j)}{||\mathbf{b}|^j - |\mathbf{a}|^j|^2}\right).$$

It follows that

$$n\mu\left(\Delta_n^{(1)} \cap T^{-j}\Delta_n\right) \leq nr_n^2 \frac{|\mathbf{a}\mathbf{b}|^{2j} \cdot |\mathbf{a}|^j}{||\mathbf{b}|^j - |\mathbf{a}|^j|^2} \leq nr_n^2 (|\mathbf{a}|^3 |\mathbf{b}|^2)^j \xrightarrow{n \rightarrow \infty} 0$$

As observed earlier, the fact that (6.8) holds implies that $p_k = 0$ for all $k \in \mathbb{N}$, which means that we are again left with computing p_0 . For that matter, note that

$$\mu\left(\Delta_n^{(1)}\right) = \mu\left(\Delta_n^{(1)} \cap I_0 \times C_0\right) + \mu\left(\Delta_n^{(1)} \cap (I_0 \times C_0)^c\right).$$

The first term is computed as in Proposition 6.5, except it comes with an extra factor of $\frac{1}{|\mathbf{a}|}$, which is precisely the measure of the set $\{x \in [0, 1] : T_1(x) = T_2(x)\}$. So, the first term is $\frac{r_n(|\mathbf{a}|-1)}{|\mathbf{a}|^2} + \mathcal{O}\left(\frac{1}{n^2}\right)$. The second term is $\frac{r_n(|\mathbf{a}|-1)}{|\mathbf{a}|} - \mathcal{O}\left(\frac{1}{n^2}\right)$. This is because the measure of $\Delta_n \cap (I_0 \times C_0)^c$ is $r_n\left(\frac{|\mathbf{a}|-1}{|\mathbf{a}|}\right) + \mathcal{O}\left(\frac{1}{n^2}\right)$ and, by Lemma 6.1 and by the fact that only a finite number of points of $D \cap (I_0 \times C_0)^c$ return to D after the first iteration, which explains that there is only a $\mathcal{O}\left(\frac{1}{n^2}\right)$ contribution. This means that the extremal index is $\theta = \frac{|\mathbf{a}|-1}{|\mathbf{a}|^2} + \frac{|\mathbf{a}|-1}{|\mathbf{a}|}$. $\square$

The proof of Proposition 6.6 provides clear insight into the mechanisms determining the value of the Extremal Index in this class of dynamics. The first contribution arises from the region where the factor maps coincide. This region represents a relative weight of $\frac{1}{|\mathbf{a}|}$ of the diagonal, and the corresponding fraction of points in Δ_n that do not return to Δ_n in one iteration converges to $\frac{|\mathbf{a}|-1}{|\mathbf{a}|}$. Their combined contribution to the Extremal Index is given by the product of these two quantities.

The complementary region, where the factor maps are misaligned, has relative weight $\frac{|\mathbf{a}|-1}{|\mathbf{a}|}$ of the diagonal. In this case, since the factor maps differ, the fraction of points in Δ_n that do not return to Δ_n converges to 1.

Equivalently, the Extremal Index can be written as $1 - \frac{1}{|\mathbf{a}|^2}$, indicating that, in the limit, a proportion $\frac{1}{|\mathbf{a}|^2}$ of points in a neighbourhood of the diagonal return to that neighbourhood after one iteration.

7. APPLICATIONS

In this section, we provide some examples to illustrate the potential of the results obtained earlier, namely Theorem 4.1 and Corollaries 5.7 and 5.8. We will introduce a class of perturbations of linear maps that fits this framework and show that when taking the direct product of perturbed versions of two dynamically incompatible maps, in the sense that the set of points that return to the diagonal is negligible, we obtain an Extremal Index equal to 1.

7.1. Perturbed linear maps. We start by recalling the definition of α -Hölder function. Let $f : I \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$. We say that f is α -Hölder if there exists $C > 0$ such that, for all $x, y \in I$

$$|f(x) - f(y)| \leq C |x - y|^\alpha. \quad (7.1)$$

It is easy to see that if f is α -Hölder and bounded away from zero (in norm), then there is $C > 0$ such that given $z \in I$, $\epsilon > 0$ and $x, y \in B_\epsilon(z)$,

$$|f(x) - f(y)| \leq C |f(z)| \epsilon^\alpha. \quad (7.2)$$

Let $f : I \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$. We say that $f \in C^{1+\alpha}(I)$ or simply that f is $C^{1+\alpha}$ if its determinant is α -Hölder.

It is clear that the product of expanding linear maps satisfies all the conditions of Theorem 4.1. Using this approach, we can guarantee that Theorem 4.1 can also be applied to small perturbations of expanding linear maps. To precisely describe what we mean by small perturbations, consider integers $\mathbf{a}, \mathbf{b}$, greater than 1 in absolute value. Let $\delta < \frac{\min\{|\mathbf{a}|, |\mathbf{b}|\} - 1}{2}$. We label the differentiable regions (open sets) of the map

$$\begin{aligned} T : [0, 1]^2 &\longrightarrow [0, 1]^2 \\ (x, y) &\longmapsto (\mathbf{a}x \bmod 1, \mathbf{b}y \bmod 1) \end{aligned}$$

by $I_{kl} = I_k \times J_l$, where $k \in \{0, \dots, |\mathbf{a}| - 1\}$, and $l \in \{0, \dots, |\mathbf{b}| - 1\}$. Consider the maps $f_k : I'_k \rightarrow \mathbb{R}$ and $g_l : J'_l \rightarrow \mathbb{R}$, where I'_k and J'_l are compact sets such that $\overline{I_k} \subseteq \text{int } I'_k$ and $\overline{J_l} \subseteq \text{int } J'_l$. Suppose that f_k and g_l are $C^{1+\alpha}$, for some $\alpha > 0$. Furthermore, assume that $f_k(\overline{I_k}) \subseteq [-1, 1]$, $g_l(\overline{J_l}) \subseteq [-1, 1]$, and that $f_k|_{\partial I_k} = 0$ and $g_l|_{\partial J_l} = 0$. Suppose also that

$$\forall x \in I'_k, y \in J'_l, |f'_k(x)|, |g'_l(y)| \leq 1.$$

Then, we consider the dynamical system

$$\tilde{T} : [0, 1]^2 \rightarrow [0, 1]^2, \tilde{T}|_{I_{kl}} = T|_{I_{kl}} + \delta(f_k|_{I_k} \times g_l|_{J_l}), \quad (7.3)$$

where

$$\begin{aligned} f_k \times g_l : I'_k \times J'_l &\mathbb{R}^2 \\ (x, y) &\longmapsto (f_k(x), g_l(y)). \end{aligned}$$

From now on, in order to simplify notation, we will write f_k instead of $f_k|_{I_k}$ and analogously for g_l . We denote $\tilde{T}_{kl}$ to the extension of $\tilde{T}|_{I_{kl}}$ to $\text{int}(I'_k \times J'_l)$. In order to apply Theorem 4.1, we check that $\tilde{T}$ satisfies (PE1)-(PE5), and that it satisfies Proposition 4.5.

First, we check that each $\tilde{T}_{kl}$ is uniformly expanding and its restriction to $\overline{I_{kl}}$ is a diffeomorphism between $\overline{I_{kl}}$ and $[0, 1]^2$. To do so, it is enough to check that each of the components are uniformly expanding and that they map, respectively, $\overline{I_k}$ and $\overline{J_l}$ to $[0, 1]$. We do it for the first component. The derivative of it is $\mathbf{a} + \delta f'_k(x)$. If $\mathbf{a} > 0$, we get $\mathbf{a} + \delta f'_k(x) > \mathbf{a}/2 + 1/2 > 1$, so the map is increasing and expanding. In particular, it is injective. By the boundary conditions imposed on the map f_k , we get that its image is $[0, 1]$. If $\mathbf{a} < 0$, the proof is similar.

The map $\tilde{T}$ satisfies condition (PE1) and (PE3), since it was actually constructed on the sets U_i (that here we label by I_{kl}) which are the same that are used in the linear setting. Condition (PE4) follows from the fact that both components are uniformly expanding. To check condition (PE2), we need the following propositions:

It remains to check $\tilde{T}$ satisfies Proposition 4.5. For that purpose, it suffices to show that for each component map of $\tilde{T}$, and for each interval $I \subseteq [0, 1]$, there exists $n \in \mathbb{N}$ such that the

image of I under that component is $[0, 1]$ (up to Lebesgue measure zero). By construction of $\tilde{T}$, there exists $\lambda \geq 1$ such that the derivative of each component map has absolute value greater than λ on each partition element. An induction argument shows that the image of any interval is either $[0, 1]$, an interval of length at least λ times the original, or a union of two intervals near 0 and 1 whose total length grows. In conclusion, Proposition 4.5 applies. Hence, Theorem 4.1 also applies.

Assuming that $|\mathbf{a}| \neq |\mathbf{b}|$ and that the maps f_k and g_l are C^2 , one can easily compute the extremal index. Let $\delta < 1/2$. Then, it is clear that for all $x \in \overline{I_k^j}$ and $y \in \overline{J_l^j}$,

$$(|\mathbf{a}| - \delta)^j \leq \left| \left(T_{i,1}^j \right)'(x) \right| \leq (|\mathbf{a}| + \delta)^j$$

$$(|\mathbf{b}| - \delta)^j \leq \left| \left(T_{i,2}^j \right)'(y) \right| \leq (|\mathbf{b}| + \delta)^j.$$

This, together with the fact that $|\mathbf{a}| \neq |\mathbf{b}|$, implies that the conditions of Corollary 5.8 are satisfied, meaning that the EVL has an extremal index equal to 1. In other words, we have just proved the following:

Theorem 7.1. *Let $\tilde{T}: [0, 1]^2 \rightarrow [0, 1]^2$ be a perturbed linear map as defined in (7.3), with $|\mathbf{a}| \neq |\mathbf{b}|$. Then, for all $s > 0$ there exists a sequence $(u_n)_{n \in \mathbb{N}}$ such that (4.3) holds and*

$$\lim_{n \rightarrow \infty} \mu(M_n \leq u_n) = e^{-s},$$

APPENDIX A. THE CO-DIMENSION C_q

Throughout the paper, we mostly considered expanding maps with smooth invariant densities bounded away from zero and infinity. In that case, the co-dimension satisfies

$$C_q = 1,$$

reflecting the fact that the measures behave locally like Lebesgue measure.

However, C_q may differ from 1 when the invariant densities exhibit singularities or zeros, or when the measures have a more irregular (e.g. fractal) structure. This occurs, for instance, in unimodal maps [12, 13].

We now give an explicit formula for C_q in the case of absolutely continuous measures whose densities have finitely many singularities or zeros.

Proposition A.1. *Let $\mu_1, \dots, \mu_q$ be probability measures on $\mathbb{R}$, absolutely continuous with respect to Lebesgue measure, with densities $h_1, \dots, h_q$. Assume that the common support*

$$I = \bigcap_{i=1}^q \text{supp}(\mu_i)$$

is a finite union of closed intervals.

Suppose that each h_i has at most finitely many singularities or zeros, located at points $s \in S \subset I$, and that near each s ,

$$h_i(x) \asymp |x - s|^{\alpha_i(s)},$$

with $-1 < \alpha_i(s) < 0$ for singularities and $\alpha_i(s) > 0$ for zeros.

Define

$$\bar{\alpha} = \min_{s \in S} \left\{ \frac{1}{q} \sum_{i=1}^q \alpha_i(s) \right\}.$$

Then:

(1) If $S = \emptyset$ or $\bar{\alpha} \geq 0$, then

$$C_q(\mu_1, \dots, \mu_q) = 1.$$

(2) If $\bar{\alpha} < 0$, then

$$C_q(\mu_1, \dots, \mu_q) = \begin{cases} 1 & \text{if } q < -\frac{1}{\bar{\alpha}}, \\ \frac{q(1 + \bar{\alpha})}{q - 1} & \text{otherwise.} \end{cases}$$

Proof. Let

$$I(r) = \int_I \prod_{i=2}^q \mu_i(B_r(x)) d\mu_1(x).$$

We split the integral into contributions near singularities and away from them:

$$I(r) = I_1(r) + I_2(r),$$

where

$$I_1(r) = \sum_{s \in S} \int_{B_r(s)} h_1(x) \prod_{i=2}^q \int_{x-r}^{x+r} h_i(y) dy dx,$$

and $I_2(r)$ is the integral over $I \setminus B_r(S)$.

For $x \notin B_r(S)$, the densities are regular, so

$$\int_{x-r}^{x+r} h_i(y) dy \asymp r.$$

Hence

$$I_2(r) \asymp r^{q-1}.$$

For x close to s , we use the local behaviour

$$\int_{x-r}^{x+r} h_i(y) dy \asymp (|x - s| + r)^{1+\alpha_i(s)}.$$

Thus

$$I_1(r) \asymp \sum_{s \in S} \int_{s-r}^{s+r} |x - s|^{\alpha_1(s)} (|x - s| + r)^{\sum_{i=2}^q (1+\alpha_i(s))} dx.$$

A direct scaling argument yields

$$I_1(r) \asymp \sum_{s \in S} r^{q + \sum_{i=1}^q \alpha_i(s)}.$$

Combining both contributions,

$$I(r) \asymp r^{q-1} + r^{q(1+\bar{\alpha})}.$$

The dominant term determines the scaling. If

$$q(1 + \bar{\alpha}) > q - 1,$$

i.e. $\bar{\alpha} > -1/q$, then $I(r) \asymp r^{q-1}$ and $C_q = 1$. Otherwise, $q(1 + \bar{\alpha}) \leq q - 1$ and the singular contribution dominates:

$$I(r) \asymp r^{q(1+\bar{\alpha})}.$$

In either case, by definition of C_q ,

$$C_q = \lim_{r \rightarrow 0} \frac{\log I(r)}{(q-1) \log r},$$

which gives $C_q = 1$ in the first case and $C_q = \frac{q(1+\bar{\alpha})}{q-1}$ in the second. $\square$

Remark A.2. The formula highlights a transition phenomenon: singularities only affect C_q when they are sufficiently strong (i.e. when $\bar{\alpha} < 0$) and when the number of trajectories q is large enough. Otherwise, the regular part of the measure dominates and $C_q = 1$.

Example A.3. Let $I = [0, 1]$.

- (1) Let $h_1(x) = h_2(x) = \frac{1}{\pi\sqrt{x(1-x)}}$ and $h_3(x) = 1$. h_1 and h_2 are the invariant densities associated with the logistic map, while h_3 is that of an expanding linear map with integer slope. Each of h_1, h_2 has exponents $\alpha_1(0) = \alpha_2(0) = \alpha_1(1) = \alpha_2(1) = -\frac{1}{2}$ and $\alpha_3 \equiv 0$, giving $\bar{\alpha} = \frac{1}{3}(-\frac{1}{2} - \frac{1}{2} + 0) = -\frac{1}{3}$. The condition $q(1 + \bar{\alpha}) > q - 1$ reads $3 \cdot \frac{2}{3} > 2$, which holds as an equality, so the two exponents coincide and

$$C_3(\mu_1, \mu_2, \mu_3) = 1.$$

- (2) Let $h_1(x) = h_2(x) = h_3(x) = h_4(x) = \frac{1}{\pi\sqrt{x(1-x)}}$ and $h_5(x) = 1$. Here $\bar{\alpha} = \frac{1}{5}(4 \cdot (-\frac{1}{2}) + 0) = -\frac{2}{5}$, and the condition $q(1 + \bar{\alpha}) > q - 1$ reads $5 \cdot \frac{3}{5} > 4$, i.e. $3 > 4$, which is false. Hence

$$C_5(\mu_1, \dots, \mu_5) = \frac{5(1 - \frac{2}{5})}{4} = \frac{3}{4}.$$

REFERENCES

- [1] R. Aimino, M. Nicol, S. Vaienti, Annealed and quenched limit theorems for random expanding dynamical systems, *Probab. Theory Related Fields* 162(1-2) (2015), 233–274.
- [2] R. Aimino, M. Nicol, M. Todd, Recurrence statistics for the space of interval exchange maps and the Teichmüller flow on the space of translation surfaces, *Ann. Inst. H. Poincaré Probab. Statist.* 53(3) (2017): 1371–1401.
- [3] J. Atnip, G. Froyland, C. González-Tokman, and S. Vaienti. Compound Poisson statistics for dynamical systems via spectral perturbation. *Nonlinearity*, 38(8):Paper No. 085023, 47, 2025.
- [4] J. Atnip, N. Haydn, and S. Vaienti. Extreme value theory with spectral techniques: application to a simple attractor. *Discrete Contin. Dyn. Syst.*, 45(10):3512–3544, 2025.
- [5] H. Aytaç, J. M. Freitas, and S. Vaienti. Laws of rare events for deterministic and random dynamical systems. *Trans. Amer. Math. Soc.*, 367(11):8229–8278, 2015.
- [6] V. Barros, L. Liao, J. Rousseau, On the shortest distance between orbits and the longest common substring problem, *Adv. Math.* 344 (2019), 311–339.
- [7] V. Barros, J. Rousseau, Shortest distance between multiple orbits and generalised fractal dimensions, *Ann. Henri Poincaré* 22(6) (2021), 1853–1885.
- [8] V. Barros, A. Coutinho, Shortest distance between observed orbits in distinct dynamical systems, *Chaos, Solitons & Fractals* 206 (2026), 117933.
- [9] T. Caby, Matching of observations of dynamical systems, with applications to sequence matching, *Phys. D.* 440 (2022) 133456.

- [10] T. Caby, D. Faranda, G. Mantica, S. Vaienti, P. Yiou, Generalized dimensions, large deviations and the distribution of rare events, *Phys. D* 400 (2019), 132–143.
- [11] T. Caby, D. Faranda, S. Vaienti, P. Yiou, On the computation of the extremal index for time series, *J. Stat. Phys.* 179 (2020), 1666–1697.
- [12] T. Caby, M. Gianfelice, *On the generalized dimensions of physical measures of chaotic flows*, *Chaos, Solitons & Fractals*, vol. 199, part 2, p. 116678, 2025.
- [13] T. Caby, M. Gianfelice, B. Saussol, S. Vaienti, *Topological Synchronization or a simple attractor*, *Nonlinearity* 36 n. 7, 3603–3621 (2023).
- [14] J.-R. Chazottes, Z. Coelho, and P. Collet. Poisson processes for subsystems of finite type in symbolic dynamics. *Stoch. Dyn.*, 9(3):393–422, 2009.
- [15] Z. Coelho and P. Collet. Asymptotic limit law for the close approach of two trajectories in expanding maps of the circle. *Probab. Theory Related Fields*, 99(2):237–250, 1994.
- [16] A. Coutinho, R. Lambert, J. Rousseau, Matching strings in encoded sequences, *Bernoulli* 26(3) (2020), 2021–2050.
- [17] A. Coutinho, J. Rousseau, B. Saussol, Large deviation for return times, *Nonlinearity* 31(11) (2018), 5162–5179.
- [18] D. Faranda, H. Ghoudi, P. Guiraud, S. Vaienti, Extreme value theory for synchronization of coupled map lattices, *Nonlinearity* 31 (2018), 26–58.
- [19] D. Faranda, S. Vaienti, Correlation dimension and phase space contraction via extreme value theory, *Chaos* 28 (2018), 041103.
- [20] D. Faranda, G. Messori, P. Yiou, Dynamical proxies of North Atlantic predictability and extremes, *Sci. Rep.* 7 (2017), 41–78.
- [21] D. Faranda, A. C. Moreira Freitas, J. M. Milhazes Freitas, M. Holland, T. Kuna, V. Lucarini, M. Nicol, M. Todd, S. Vaienti, *Extremes and Recurrence in Dynamical Systems*, Wiley, New York (2016).
- [22] A. C. M. Freitas, J. Freitas, M. Todd, Hitting Time Statistics and Extreme Value Theory, *Probab. Theory Related Fields* 147(3) (2010), 675–710.
- [23] A. C. M. Freitas, J. M. Freitas, On the link between dependence and independence in extreme value theory for dynamical systems, *Statist. Probab. Lett.* 78(9) (2008), 1088–1093.
- [24] A. C. M. Freitas, J. M. Freitas, M. Todd, Speed of convergence for laws of rare events and escape rates, *Stochastic Process. Appl.* 125(4) (2015), 1653–1687.
- [25] A. C. M. Freitas, J. M. Freitas, J. V. Soares, Rare events for product fractal sets, *J. Phys. A: Math. Theor.* 54 (2021), 345202.
- [26] M. Carvalho, A. C. M. Freitas, J. M. Freitas, M. Holland, M. Nicol, Extremal dichotomy for uniformly hyperbolic systems, *Dyn. Syst.* 30(4) (2015), 383–403.
- [27] S. Gouëzel, P. Rousseau, M. Stadlbauer, Minimal distance between random orbits, *Probability Theory and Related Fields* 189 (2024), 811–847.
- [28] G. Keller, Rare events, exponential hitting times and extremal indices via spectral perturbation, *Dyn. Syst.* 27(1) (2012), 11–27.
- [29] G. Keller, C. Liverani, Stability of the spectrum for transfer operators, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) 28(1) (1999), 141–152.
- [30] G. Keller, C. Liverani, Rare events, escape rates and quasistationarity: some exact formulae, *J. Stat. Phys.* 135 (2009), 519–534.
- [31] J. Liu, S. Shi, Shortest distance between partial orbits, *Discrete and Continuous Dynamical Systems – Series S*, to appear.
- [32] J. Rousseau, M. Todd, Orbits closeness for slowly mixing dynamical systems, *Ergod. Th. & Dynam. Sys.* 19 (2023), 1–17.
- [33] B. Saussol, Absolutely continuous invariant measures for multidimensional expanding maps, *Israel J. Math.* 116 (2000), 223–248.
- [34] S. Shi, J. Liu, and Y. Zhang. The shortest distance between orbits of two different dynamical systems. *Acta Math. Hungar.*, 2026. (DOI: 10.1007/s10474-026-01596-w)
- [35] S. Shi, On the shortest distance between orbits in conformal iterated function systems, preprint.

ROMAIN AIMINO, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL

Email address: `romain.aimino@fc.up.pt`

URL: `https://sites.google.com/view/romain-aimino/accueil`

THÉOPHILE CABY, LABORATOIRE D'Océanographie Physique et Spatiale, UNIV BREST CNRS IFREMER IRD, BREST, FRANCE,

Email address: `theophile.caby@uni-brest.fr`

JORGE MILHAZES FREITAS, CENTRO DE MATEMÁTICA & FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL

Email address, Corresponding author: `jmfreira@fc.up.pt`

URL: `http://www.fc.up.pt/pessoas/jmfreira/`

DUARTE SÁ PINHO, CENTRO DE MATEMÁTICA & FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL

Email address: `duarte.sa.pinho02@gmail.com`